

Zeta Functions from Definable Equivalence Relations

Ehud Hrushovski and Ben Martin

February 2, 2008

Abstract

We prove that the theory of the p -adics $\mathbb{Q}_p$, together with a set of explicitly given sorts, admits elimination of imaginaries. Using p -adic integration, we deduce the rationality of certain formal zeta functions arising from definable equivalence relations. As an application, we prove rationality results for zeta functions obtained by counting isomorphism classes of irreducible representations of finitely generated nilpotent groups; these are analogous to similar results of Grunewald, Segal and Smith for subgroup growth zeta functions of finitely generated nilpotent groups.

1 Introduction

[Warning: This is a preliminary version. There are inconsistencies of notation, missing references and citations, and probably some minor mathematical mistakes and gaps.]

This paper concerns the model theory of the p -adic numbers $\mathbb{Q}_p$ and applications to certain counting problems arising in group theory. Recall that a theory (in the model-theoretic sense of the word) is said to have *elimination of imaginaries* (EI) if the following holds: for every model M of the theory, for every definable subset D of some M^n and for every definable equivalence relation R on D , there exists a definable function $f: D \rightarrow M^m$ for some m such that the fibres of f over $f(D)$ are precisely the equivalence classes of R . The theory of $\mathbb{Q}_p$ (in the language of valued rings) does not admit EI: for example, no such f exists for the definable equivalence relation

R on $\mathbb{Q}_p$ given by xRy if $|x - y| \leq 1$, because $\mathbb{Q}_p/R$ is countably infinite but any definable subset of $\mathbb{Q}_p^m$ is either finite or uncountable.

We show that the theory of $\mathbb{Q}_p$ plus some extra sorts admits EI. In fact, we prove a more general result (Proposition 2.4): given two theories $T, \tilde{T}$ satisfying certain hypotheses, T has EI if $\tilde{T}$ does. In our application, $\tilde{T}$ is the theory $ACVF_{0,p}$ of algebraically closed fields of characteristic zero with residue field of characteristic p and T is the theory of $\mathbb{Q}_p$, with appropriate extra sorts in both cases.

The notion of an invariant extension of a type plays a key part in our proof. If T is a theory, $M \models T$, $A \subseteq M$ and p is a type over A then an *invariant extension* of p is a type q over M such that $q|_A = p$ and q is $\text{Aut}(M/A)$ -invariant. The theory $ACVF_{0,p}$ is not stable; in [5], [6], Haskell, the first author and Macpherson used invariant extensions of types to study the stability properties of $ACVF_{0,p}$ and to define notions of forking and independence. They proved that $ACVF_{0,p}$ plus some extra sorts admits EI. We use not only this result but also other properties of types given in [5], [6] in our proof of p -adic elimination of imaginaries.

Our main application of p -adic EI is to rationality results for certain zeta functions associated to finitely generated nilpotent groups. Grunewald, Segal and Smith [4] showed that subgroups of p -power index of such a group Γ can be parametrised p -adically. More precisely, these subgroups can be *coded*: that is, placed in bijective correspondence with the set of equivalence classes of some definable equivalence relation on a definable subset D of some $\mathbb{Q}_p^N$. Let $a_n < \infty$ denote the number of subgroups of Γ of index n . Using p -adic integration over D and results of [Igusa, Denef, etc.], Grunewald, Segal and Smith showed that the p -local subgroup growth zeta function $\sum_{n=0}^{\infty} a_{p^n} t^n$ is a rational function of t . Du Sautoy and his collaborators have calculated subgroup growth functions explicitly in many cases, and studied uniformity questions (the behaviour of the p -local subgroup growth function as the prime p varies).

A crucial step in using p -adic integration to prove rationality is to show that there exists a definable function $f: D \rightarrow \mathbb{Q}_p$ such that the Haar measure of each equivalence class $[x]_R \in D/R$ of the form

$$\mu([x]_R) = |f(x)|; \tag{1}$$

in practice, f is usually given explicitly (cf. [formulas from [4]]). This is straightforward for the case of p -power index subgroups of Γ coded in the

usual way. For more complicated equivalence relations, however, it is not clear that such an f can be found, even in principle.

A consequence of elimination of imaginaries is that one can reduce an arbitrary definable equivalence relation to a definable equivalence relation of a particular kind — namely, the equivalence relation on $GL_N(\mathbb{Q}_p)$ for some N whose equivalence classes are the left $GL_N(\mathbb{Z}_p)$ -cosets. We give a formula (Eqn. 2) of the form (1) for the Haar measure of the equivalence classes. Using results on p -adic integration due to Denef, we prove Theorem 6.2, which says that zeta functions that arise from counting the number of equivalence classes of certain definable equivalence relations are rational.

We expect Theorem 6.2 will be a useful tool for proving rationality results. We give some applications to situations in which it is not clear how to apply the definable p -adic integral formalism. The main one, and the original motivation for our results, is in the area of representation growth. This is analogous to subgroup growth: one counts not the number b_{p^n} of index p^n subgroups of a group Γ , but the number a_{p^n} of irreducible p^n -dimensional complex characters of Γ (modulo tensoring by one-dimensional characters if Γ is nilpotent). Representation growth of finitely generated pro- p groups was studied by Jaikin [refs]; Lubotzky and the second author gave a partial criterion for an arithmetic group to have the congruence subgroup property in terms of its representation growth [9, Theorem 1.2]. We prove that the p -local representation growth zeta function $\zeta_{\Gamma,p}(t) = \sum_{n=0}^{\infty} a_{p^n} t^n$ of a finitely generated nilpotent group Γ is rational (Theorem 8.4). We hope that Theorem 8.4 will stimulate further investigation of representation growth, as the paper [4] did for subgroup growth.

Jaikin [ref] has proved, under mild technical restrictions, that the p -local representation growth functions of semisimple compact p -adic analytic groups are rational; we are grateful to him for explaining his work. His parametrisation of irreducible characters uses the coadjoint orbit formalism of Howe; rationality follows from the usual methods of p -adic integration. [More explicit: e.g. SL_2 ?] We believe that his methods could be adapted to prove rationality in the case of finitely generated nilpotent groups as well.

This paper falls naturally into two parts. The first part is model-theoretic: we establish our main results Proposition 2.4 and Theorem 2.2 on elimination of imaginaries, and establish the general rationality result Theorem 6.2. In the second part, we apply Theorem 6.2 to prove rationality of some group-theoretic zeta functions, including the representation growth zeta functions for finitely generated nilpotent groups (Theorem 8.4). The main tools are

results from profinite groups; no ideas from model theory are used in a significant way beyond the notion of definability.

The authors wish to thank Thomas Rohwer, Deirdre Haskell and Dugald Macpherson for their comments on earlier drafts of this work. We are grateful to Alex Lubotzky for suggesting studying representation growth; several of the ideas in Section 8 are due to him. The second author was supported by a Golda Meir Postdoctoral Fellowship at the Hebrew University of Jerusalem.

2 Elimination of Imaginaries

2.1 Model-Theoretic Preliminaries

Let E be a definable relation, that is, a definable subset of M^n for some n . Given a formula $R(x, y)$, where x is an n -tuple of variables, and given a tuple b , we denote by $R(x, b)$ the set $\{x \in M^n : M \models R(x, b)\}$.

A definable relation E is *coded* if it can be written as $R(x, b)$, where b is a tuple of elements of the sorts of the theory, and where $b \neq b'$ implies that $R(x, b)$ does not coincide with $R(x, b')$. In this situation b is called a *code* for E ; $dcl(b)$ depends only on E , and is denoted $\langle E \rangle$. T is said to eliminate imaginaries (EI) if every definable relation in every model of T is coded. ACVF has EI for the sorts S_n, T_n plus the dominant field sort K ([HHM]).

In general, we use the notation $\langle E \rangle$ to refer to the code of E with respect to T^{eq} .

2.2 Many-Sorted Structures

We will consider many-sorted theories with a distinguished collection $\mathcal{S}$ of sorts, referred to as the *dominant* sorts; we assume that for any sort S , there exists a 0-definable partial function from a finite product of dominant sorts onto S (and this function is viewed as part of the presentation of the theory.)

The set of elements of dominant sorts in a model M is denoted $\text{dom}(M)$.

Lemma 2.1 (Cf. [5, Remark 3.2.2]) *A theory T admits elimination of imaginaries if every definable function on a dominant sort, in any model of T , is coded.*

Proof It suffices to show that every definable function f on M^n ($n \in \mathbb{N}$) is coded. Pulling back by the given 0-definable functions, it suffices to show that every definable function on a product $M_1 \times \dots \times M_n$ of dominant sorts is coded. For $n = 1$, this is our assumption. For larger n , we use induction, regarding a definable function from $M_1 \times \dots \times M_n$ to M as a definable function from M_n into the set of definable functions from $M_1 \times \dots \times M_{n-1}$ to M , or rather, into the set of codes for these functions.

2.3 Valued fields

Let K be a valued field, with valuation ring $\mathcal{O}$, residue field k , value group Γ , maximal ideal $\mathcal{M}$.

We take a single dominant sort, for K itself. The additional sorts S_n, T_n for $n \in \mathbb{N}$ are given by

$$S_n(K) = GL_n(K)/GL_n(\mathcal{O}) = B_n(K)/B_n(\mathcal{O}),$$

the set of lattices in K^n , and

$$T_n(K) = GL_n(K)/GL_{n,n}(\mathcal{O}) = \bigcup_{m \leq n} B_n(K)/B_{n,m}(\mathcal{O}) = \bigcup_{e \in S_n} e/\mathcal{M}e.$$

Here a *lattice* is a free $\mathcal{O}$ -submodule of K^n of rank n , B_n is the group of upper triangular matrices, $GL_{n,m}(\mathcal{O})$ is the group of matrices in $GL_n(\mathcal{O})$ whose m th column reduces mod $\mathcal{M}$ to the column vector $(1, 0, 0, \dots, 0)$ of k , and $B_{n,m}(\mathcal{O}) := B_n(\mathcal{O}) \cap GL_{n,m}(\mathcal{O})$. There is a canonical map from $T_n(K)$ to $S_n(K)$ taking $f = e/\mathcal{M}e$ to the lattice $\mathcal{M}$.

It is easy to see using elementary matrices that $GL_n(K) = B_n(K)GL_n(\mathcal{O})$, justifying the equality of the first two definitions of S_n . Equivalently, it is shown in [5, Lemma 2.4.8] that every lattice has a basis in triangular form.

Note that k can be identified with the fiber of $T_1 \rightarrow S_1$ above 0, while the value group Γ can be identified with S_1 . More generally, if $B \subseteq K$ then let $\mathbb{B}(B) = \{\{x : |x - a| \leq |b|\} : a, b \in B\}$. Then $\mathbb{B}$ embeds into $S_2 \cup K$. (The balls of 0-radius are identified with K . The group G of affine transformations of the line acts on $\mathbb{B}$ transitively on the balls of nonzero radius; the stabilizer of $\mathcal{O} \in \mathbb{B}$ is $G(\mathcal{O})$; so $\mathbb{B} \setminus K \simeq G(K)/G(\mathcal{O}) \leq GL_2(K)/GL_2(\mathcal{O})$.)

The theory of a structure is determined by the theory of the dominant sorts; so we can speak e.g. of $Th(\mathbb{Q}_p)$ in these sorts. We take the language to include the ring structure on K , and the natural maps $GL_n(K) \rightarrow S_n(K)$, $GL_n(K) \rightarrow T_n(K)$.

Theorem 2.2 *The theory of $\mathbb{Q}_p$ in the field sort together with the sorts S_n eliminates imaginaries.*

We give a proof of this using the theory of algebraically closed valued fields, and invariant types. Let us say that a theory T has the invariant extension property IE (respectively has property IE for a sort S) if whenever $A = \text{acl}(A) \subseteq M \models T$, and $c \in M$ (respectively $c \in S$), $\text{tp}(c/A)$ extends to an $\text{Aut}(M/A)$ -invariant type over M . This holds trivially for any finite field, and by inspection, for $\text{Th}(\mathbb{Z}, +, <)$. We will show that it holds for the p -adics too.

2.4 Real elimination of imaginaries

To illustrate the idea, consider this way of deducing EI for RCF (the theory of real closed fields) from EI for ACF (the theory of algebraically closed fields). Later we will see how to replace ACF by any stable theory: indeed any theory with enough invariant types of the right kind.

Example 2.3 Let F be a field. Assume:

- (i) The theory of F is algebraically bounded;
- (ii) (Rigidity of finite sets) No automorphism of a model of T can have a finite cycle of size > 1 . Equivalently, for each n , $\text{Th}(F)$ has symmetric 0-definable functions $r_{i,n}(x_1, \dots, x_n)$, such that (denoting $r_{i,n}(S) = r_{i,n}(x_1, \dots, x_n)$ when $S = \{x_1, \dots, x_n\}$), $S = \{r_{1,n}(S), \dots, r_{n,n}(S)\}$;
- (iii) (Unary EI) Every definable subset of F is coded.

Then F eliminates imaginaries (in the single sort of field elements.)

Proof Let f be a definable function on F . By Lemma 2.1, it suffices to prove that f is coded. Let H be the Zariski closure of the graph of f . Since the theory is algebraically bounded, the set $H(x) := \{y : (x, y) \in H\}$ is finite for any x , of size bounded by some n . Let $U_{n,i}$ be the set of x such that $f(x) = r_{i,n}(H(x))$. Then H (being an F -Zariski closed set) and each U_i (being unary) is canonically coded; these codes together give a code for f .

Note that RCF satisfies the hypotheses of Example 2.3 [check!], but $\text{Th}(\mathbb{Q}_p)$ (in the field sort alone) does not. [Justify: $\text{Th}(\mathbb{Q}_p)$ doesn't have unary EI.]

2.5 Criterion for Elimination of Imaginaries

Proposition 2.4 *Let $\tilde{L} \subseteq L$ be languages. Let T be a complete L -theory, $\tilde{T}$ a complete $\tilde{L}$ -theory admitting elimination of quantifiers and of imaginaries, and assume T contains the universal part of $\tilde{T}$. Then T too admits elimination of imaginaries, provided conditions (i)-(v) below hold.*

The closure under quantifier-free $\tilde{L}$ - definable functions is denoted dcl ; the closure under L -definable functions in $M \models T$ is denoted DCL . Similarly acl , ACL and tp, TP .

Assume that whenever M is a sufficiently saturated model of T , $A = ACL(A) \subseteq M$, $c \in \text{dom}(M)$, $M|_{\tilde{L}} \leq \tilde{M} \models \tilde{T}$, we have:

- (i) *(Relative algebraic boundedness.) For every elementary submodel M' of M , $DCL(M'c) \subseteq acl(M'c)$.*
- (ii) *(Weak rigidity). $ACL(Ac) \subseteq DCL(Ac)$.*
- (iii) *(Unary EI) Every M -definable subset of $\text{dom}(M)$ is coded in M (or just by elements of $\tilde{M}$ definable over M .)*
- (iv) *(Internalizing $\tilde{L}$ -codes.) Let $e \in \tilde{M}$ and suppose $e \in dcl(M)$. Then there exists a tuple e' of elements of M such that an automorphism of $\tilde{M}$ leaving M setwise invariant fixes e iff it fixes e' .*
- (v) *(Invariant types with strong germs.) There exists an $\text{Aut}(\tilde{M}/A)$ -invariant type $\tilde{p}$ over $\tilde{M}$ such that $\tilde{p}|_M$ is consistent with $TP(a/A)$.*
Moreover, for any $\tilde{L}$ -definable function r on $\tilde{M}$, let $\partial_{\tilde{p}}r$ be the $\tilde{p}$ -germ of r (where two functions r, r' defined over $\tilde{M}$ have the same $\tilde{p}$ -germ if they agree on a realization of $\tilde{p}$ over $\tilde{M}$.) Then:
 - (*) *There exists a sequence $e_i \in dcl(A, \langle r \rangle)$ such that $\sigma \in \text{Aut}(\tilde{M}/A)$ fixes $\partial_{\tilde{p}}r$ iff σ fixes almost every e_i .*
 - (**) *If $\partial_{\tilde{p}}r$ is fixed by $\text{Aut}(\tilde{M}/A)$, then there exists an A -definable function h (for $\tilde{L}, \tilde{T}$) with the same $\tilde{p}$ -germ.*

Proof Let M be a rather saturated and homogeneous model of T . By Lemma 2.1, it suffices to show that M -definable functions on $\text{dom}(M)$ are coded. Let f be such a function. Let $A = ACL(\langle f \rangle) \cap M$ and let $A' = DCL(A, \langle f \rangle)$ (allow A' to include imaginary elements.)

Claim 2.5 There exists a binary relation R , $\tilde{L}$ -definable over M , such that for any c , $R(c) := \{a : \tilde{M} \models R(c, a)\}$ is a finite set, and for $c \in \text{dom}(M)$, $f(c) \in R(c)$.

Proof By relative algebraic boundedness (i), and compactness.

Claim 2.6 For any complete type $p = tp(c/A)$ ($c \in \text{dom}(M)$), f agrees with some A -definable function on p .

Proof Let $\tilde{p}$ be a type as in (v), with the “strong germ” property (*), (**). For any R as in Claim 2.5, there exists some finite $m = m(R)$ such that if $c \models \tilde{p}|M$ then $R(c)$ is an m -element set. Let m_0 be the minimal possible value of $m(R)$, and restrict attention to relations R satisfying the conclusion of Claim 2.5, such that $m(R) = m_0$. If R, R' are two such relations, then their conjunction has the same property. But as $m(R \& R') \leq m(R) = m(R') = m_0$, equality must hold. It follows that for $c \models \tilde{p}|M$, $R(c) = R'(c)$. Let $r(c)$ be a code for the finite set $R(c)$. The $\tilde{p}$ -germ $\langle \partial_{\tilde{p}} r \rangle$ of r is fixed by $\text{Aut}(M/A')$. Let (e_i) be a sequence as in (v) (*). Note that $e_i \in \text{dcl}(M)$, so it makes sense to talk of $\text{Aut}(M)$ acting on the e_i . Let e_i' be as in (iv). By Neumann’s Lemma, all but finitely many e_i must be $\text{Aut}(M/A')$ -invariant. Then all but finitely many e_i' must be in $\text{ACL}(A')$. (If an infinite set $e_i', i \in J$ lies outside $\text{ACL}(A')$, some automorphism $\tau \in \text{Aut}(M/\text{ACL}(A'))$ would have $\tau(e_j) \neq e_j$ for $j \in J$, but this contradicts that τ fixes the germ.) But e_i' are real elements, so $e_i' \in A$. Let $A^* = \{e \in \text{dcl}(M) : \text{Aut}(M/A)e = e\}$. Then $e_i \in A^*$ for each such i , so $\text{Aut}(\tilde{M}/A^*)$ fixes each e_i , and by (v) (*), the $\tilde{p}$ -germ of r is invariant under $\text{Aut}(\tilde{M}/A^*)$. By (**), there exists an A^* -definable function r' with the same $\tilde{p}$ -germ as r .

Let $R'(x)$ be the finite set coded by $r'(x)$, $R''(x) = R'(x) \cap M$. Then $R''(x)$ is $\text{Aut}(M/A)$ -invariant, hence an A -definable function in M . Let $c \models \tilde{p}|B$ (where B contains A and f is defined over B .) By (ii), $R''(c) \subseteq \text{DCL}(Ac)$. As $f(c) \in R(c) = R'(c)$, we have $f(c) \in R''(c)$, so $f(c) = g(c)$ for some A -definable g .

Let $E = \{x : f(x) = g(x)\}$. Then E is A' -definable. Using unary EI (iii), E has a code e ; clearly e is defined almost over $\langle f \rangle$, so $e \in A$, and thus E is A -definable. As $f(c) = g(c)$ when $c \models p$, we have $E \in p$. The claim follows.

Claim 2.7 The function f is A -definable.

Proof By Claim 2.6 and compactness, there exist A -definable sets $D_1, \dots, D_m$ whose union covers $\text{dom}(f)$, and A -definable functions F_i on D_i agreeing with $f|D_i$; so $f = \cup_i F_i$ is A -definable.

Now to prove the proposition, let a be a tuple from A , such that f is a -definable. Let e be the (finite) set of conjugates of a over $\langle f \rangle$. Then $DCL(\langle f \rangle) = DCL(e)$. Now by EI for $\tilde{T}$, the finite set e is coded by a tuple e' , though e' may consist of elements in $dcl(M)$ but outside M . However by (iv), there exists a tuple e'' of elements of M such that an automorphism of $dcl(M)$ leaving M invariant fixes e'' iff it fixes e' . Thus an automorphism of M fixes $\langle f \rangle$ iff it fixes e'' ; as M is finitely homogeneous, $DCL(\langle f \rangle) = DCL(e'')$, so f is coded.

Porism 2.8 The assumption $(**)$ in (v) is a little too strong for the intended application. However, it is only used for the function r of Claim 2.6; we will show that it holds for some such r .

Lemma 2.9 *Assume that for any $a \in M$, there exists a tuple c from $\text{dom}(M)$ with $a \in DCL(c)$ and such that $TP(c/ACL(a))$ extends to an $Aut(M/ACL(a))$ -invariant type over M . Then (ii) follows from:*

(ii') *If $B \subseteq \text{dom}(M)$, $acl(B) \cap M \subseteq DCL(B)$.*

Proof Let $A = \{a_i : i < \kappa\}$. For each i , pick a tuple c_i of elements of $\text{dom}(M)$ with $a_i \in DCL(c_i)$, and extend $TP(c_i/A)$ to an $Aut(\tilde{M}/A)$ -invariant type p_i . Let $A_0 = A$, and recursively let $A_{i+1} = A_i \cup \{a_i\}$, where $a_i \models p|A_i \cup \{c\}$, and $A_\lambda = \cup_{i < \lambda} A_i$ for limit λ .

Claim 2.10 $ACL(Ac) \cap DCL(A_i c) \subseteq DCL(Ac)$

Proof By induction on i . The limit case is trivial. To move from i to $i+1$, just note that as $tp(a_i/ACL(A_i c))$ does not split over A_i , any permutation of $ACL(Ac)$ that is elementary over A_i must remain elementary over A_{i+1} .

Now $DCL(A_\lambda, c) = DCL(A_\lambda \cap \text{dom}(M), c)$, so by (ii') this set contains $ACL(A_\lambda, c) \cap M$ and hence $ACL(Ac) \cap M$. Applying Claim 2.10 with $i = \lambda$, we obtain (ii).

3 The p -adic Case

We verify the hypotheses of Proposition 2.4–Porism 2.8 for $T = Th(\mathbb{Q}_p)$, $\tilde{T} = ACVF_{0,p}$.

(i) In fact, in this case, by model completeness of $Th(\mathbb{Q}_p)$ and the nature of the axioms, $acl(M'c) \cap M$ is an elementary submodel of M .

(ii) We use Lemma 2.9: the hypothesis of Lemma 2.9 is proved in Corollary 4.7, while the assumption (ii') of Lemma 2.9 is in van den Dries's thesis: cf. [11].

(iii) (P. Scowcroft has proved a weak version of this, where the sets are classes of equivalence relations in two variables. Raf Cluckers has suggested that a strong version may be true, where the codes are themselves in $\mathbb{B}$.)

Here is a proof using the present ideas. Let e be an imaginary code for a subset $D \subseteq \text{dom}(M)$. Let $A = ACL(e) \cap \mathbb{B}$, $B = ACL(e)$.

Claim 3.1 For all $c \in \text{dom}(M)$, $tp(c/A) \Rightarrow tp(c/Ae)$.

Proof Recall the description in Lemma 4.5 (cf. also Corollary 4.7) of $tp(c/A)$ and $tp(c/B)$. If $w \in \mathbb{B}(B)$ then $w \in A$, by definition of A . Thus $\cap W(c; B) = \cap W(c; A)$. Moreover $\Gamma \cap DCL(B) = \Gamma \cap DCL(A)$ (as elements of Γ are coded by balls.) Thus $tp(c/A)$ and $tp(c/B)$ are generated by exactly the same set of formulas.

Thus some 1-types over A imply D , and all the others imply $\neg D$, so that by compactness D is definable over A . Hence $e \in A$. The proof is concluded as was Proposition 2.4: there is a tuple a from $\mathbb{B}$ with $a \in ACL(e)$ and $e \in DCL(a)$; so $DCL(e) = DCL(F)$, where F is the finite set of conjugates of a over e . We already know that finite sets are coded (e.g., by Proposition 2.4 (iv) and EI in ACVF.)

(iv) Any element e of $S_n(M)$ has a basis in some finite extension L_0 of $K = \text{dom}(M)$. Say $[L_0 : K] = m_0$; let L be the join of all field extensions of K of degree m_0 . Then L is a finite extension of K ; $[L : K] = m$. (Here we use the fact that the Galois group of the p -adics is finitely generated; for more general fields, we would require canonical data for finite Galois extensions; but L is 0-interpretable and requires no code.) Moreover as $\mathbb{Q}_p \cap \tilde{\mathbb{Q}}$ is an elementary submodel of K , one can find a generator a of L over K , whose

characteristic polynomial is over $DCL(\emptyset)$ in M . One has an a -definable isomorphism $f_a : L \rightarrow K^m$ (as vector spaces over K), with $f_a(\mathcal{O}_L) = \mathcal{O}^m$. (I.e. $\mathcal{O}_L$ is a free $\mathcal{O}$ -module of rank m .) f_a further induces an isomorphism of the lattice $e \cap L^n$ with a lattice e' of K^{mn} ; $e' = f_a(e)$. As e', e are defined over L and are $Aut(L/K)$ -invariant (e' is actually defined over K), we see that if a' is $Aut(L/K)$ -conjugate to a , then $e' = f_{a'}(e)$ as well. Thus $e' = f(e)$ for any of the (finitely many) conjugates of f_a , so $DCL(e) = DCL(e')$ in the sense required in (iv). Similarly for T_n (alternatively, for finite extensions K' of K , as the value group has a least element, elements of $T_n(K')$ can be coded by elements of $S_n(K')$; cf. the proof of Theorem 2.2 below.)

(v) If a ball e is in $acl(A)$, and $c \in e$, let γ be the least radius in $\Gamma(M)$ with $\gamma \geq rad(c)$; let e_1 be the ball around e of radius γ . Then $c \in e_1$, and $rad(e_1) \in M$, so $e_1 \in M$. Also $e_1 \in ACL(A)$ so $e_1 \in A$. Every sub-ball of e_1 whose radius is finitely many notches less than γ must also be in A (since there are only finitely many such sub-balls.) In particular, some ball containing c of radius less than γ is defined over A . We conclude that $tp(e/acl(A))$ is the generic type of a strictly infinite intersection P of balls; and that P is already the intersection of a sequence of M -balls, i.e., balls with a point in M and radius in M . We know that the generic type of P extends to an $Aut(\tilde{M}/A)$ -invariant type, α_P .

Now let A, r, f be as in Claims 2.5, 2.6 of the proof of Proposition 2.4: f is an A -definable function in M . With $A' = \{d \in M^{eq} : d \in DCL(A, \langle f \rangle)\}$ $A = A' \cap M$. r is a function definable over $\tilde{M}$, with $r(t)$ an n -element set, and $f(t) \in r(t)$, for all $t \in P$ outside some proper sub-ball E_r . We minimized n , and saw that the P -germ of r is $Aut(\tilde{M}/A)$ -invariant.

We have $r(t) = \cup_{i=1}^l r_i(t)$, where r_i is irreducible on the generic type of P (cf. Definition 5.1 for irreducibility, and for βP .) r_i can be taken to be defined over M^{alg} ; by taking the intersection of conjugates over M , we can assume it is defined, in $\tilde{M}$, with parameters from M .

We have $f(t) \in r_i(t)$ for $t \in X_i$; where the X_i are M -definable sets, and $P(M) \setminus E_r \subseteq \cup_i X_i$. We may assume that no X_i is contained in a proper sub-ball of P . (Else delete it and increase E_r .)

For $b \in \beta P|A$, $r_i|b$ has a germ, coded say by $c(b) = \partial_b r_i$. Let c denote the map $b \mapsto \partial_b r$. Denote $A'' = ACL(A, \langle c \rangle, \langle r \rangle, \langle f \rangle, d) \cap M$, where d is some element of P . Let $Y_i = \{|d - t| : t \in X_i\}$. Observe that (as $\Gamma(M)$ is stably embedded in M , and the induced structure is that of Presburger arithmetic) Y_i contains an arithmetic progression restricted to $rad(P|A'')$;

thus for some finite m , any interval in $rad(P|A'')$ of length at least m must include a point of Y_i . Let us call an interval of Γ *short* if it is contained in one of the form $[s\lambda_0^n, s]$ for some $n \in \mathbb{N}$.

Claim 3.2 Let $c', c'' \in Aut(M/A')c$. For all $b \in \beta_P|A''$, if $rad(b) \in Y_i$ then $c'(b) = c''(b)$.

Proof If $r', r'' \in Aut(M/A')r$ are conjugates of r_i , so that $c'(b) = \partial_b r'$, $c''(b) = \partial_b r''$, we have to show that r', r'' have the same germ on α_b .

Otherwise, one of them, say $\partial_b r'$, differs from $\partial_b r$. So for all $t \in b$, except perhaps for a some finite union of proper sub-balls, $r_i(t) \not\subseteq r'(t)$. As r_i is irreducible on P , by Lemma 5.3, $r_i|b$ is irreducible on α_b . So $r_i(t) \cap r'(t) = \emptyset$ for $t \in b$, away from a finite union e of proper sub-balls. But there exists $t \in X_i$ with $|t - d| = rad(b)$. By completeness of $\alpha_P|A''$, and $\beta_P|A''$, $t \in b$, and $t \notin e$. However $f(t) \in r'(t)$, and also $f(t) \in r_i(t)$. This contradiction proves the claim.

Claim 3.3 There exists $\gamma_0 \in \Gamma(A)$ and an A -definable balls b_0, P_1 (or $b_0 = \emptyset$) with $b_0 \subset P \subset P_1$, such that for any $c', c'' \in Aut(M/A')c$, for some finite union of short intervals $U(c', c'')$, if $b_0 \subseteq b \subset P_1$, $rad(b) \geq \gamma_0$, and $rad(b) \notin U(c', c'')$, then $c'(b) = c''(b)$.

Proof Let $U_P(c', c'') = \{rad(b) : b \in \beta_P|A'', c'(b) \neq c''(b)\}$. By Claim 3.2, $U_P(c', c'')$ contains no point of Y_i . Now $U_P(c, c')$ is a finite union of intervals; they must each be short, by the remark regarding Y_i . By compactness, some γ_0, b_0 over A'' , for all c', c'' , for some finite union $U(c, c')$ of short intervals, for all b , if $b_0 \subseteq b \subset P$, $rad(b) \geq \gamma_0$, and $rad(b) \notin U(c, c')$ then $c'(b) = c''(b)$. Since such γ_0, b_0 , exist, one can take γ_0 over A . If P has an A -definable sub-ball, one can take b_0 to be that sub-ball. If not, then all sub-balls of P of a given radius have the same type over A' ; since some b_0 has the stated property, they all do; but any sufficiently large ball b contains a ball b_0 of the prescribed size; so in that case b_0 is irrelevant, and can be taken to be $\emptyset$. Finally, use compactness to replace P by P_1 .

Claim 3.4 Let $c_\Gamma = DCL(A', <c>) \cap \Gamma$. Let $c', c'' \in Aut(M/A'c_\Gamma)c$. Then $c'(b) = c''(b)$ for all $b \in \beta_P|A$ with $rad(b) \notin c_\Gamma$.

Proof Let

$$W(c', c'') = \{a \in \Gamma(M) \setminus c_\Gamma : (\exists b)(b_0 \subseteq b \subset P_1, rad(b) \geq \gamma_0, c(b) \neq c'(b), \lambda_0 a \leq rad(b) \leq a)\}$$

I.e. $W(c', c'')$ is the set of left endpoints of "unit intervals" meeting $U(c', c'')$. By Claim 3.3, $W(c', c'')$ is finite. Clearly $W(c_1, c_3) \subseteq W(c_1, c_2) \cup W(c_2, c_3)$. Moreover, as $c' \in \text{Aut}(M/A'_{c_\Gamma})c$, $ACL(A'_{c_\Gamma}c') \cap \Gamma = DCL(A'_{c_\Gamma}c') \cap \Gamma = DCL(A'_{c_\Gamma}c) \cap \Gamma = DCL(A'c) \cap \Gamma = c_\Gamma$, so that $ACL(A'_{c_\Gamma}c') \cap (\Gamma \setminus c_\Gamma) = \emptyset$. Thus the hypotheses of Lemma 5.5 apply to $\text{Aut}(M/A'_{c_\Gamma})c, \Gamma \setminus c_\Gamma, U$. By Lemma 5.5, $W(c_1, c_2) = \emptyset$. The claim follows.

Claim 3.5 There exists an A'_{c_Γ} -definable function d_1 (for $\tilde{M}$), and an A'^{c_Γ} -definable set Q containing $\{b \in \beta_P|A : \text{rad}(b) \notin c_\Gamma\}$, and an A'_{c_Γ} -definable function d_1 on Q , such that $d_1(b) = \partial_b(r|b)$.

Proof Follows from Claim 3.4 and compactness.

Claim 3.6 There exist $\tilde{M}$ -definable functions $d, \tilde{r}$, with d defined over A , such that for $b \in \beta_P|A$,

$$d(b) = \partial_b(\tilde{r}|b) \text{ and } f(t) \in \tilde{r}(t). \quad (*)$$

Proof $(*)$ holds for $\tilde{r} = r$ and d_1 , for $b \in Q$, by Claim 3.5; moreover we may write $d_1(u) = d_1(\bar{\gamma}, u)$, $Q = Q(\bar{\gamma}, t)$, with $\bar{\gamma}$ certain parameters from $c_\Gamma \subset \Gamma$. Further, we can take Q to contain all elements of $\beta_P|A$ with radius outside $\bar{\gamma}$. As Γ is Skolemized over A' , we can find $\bar{\gamma}'$ from $\Gamma(A')$ such that for some $\tilde{r}$, $(*)$ holds true for all $b \in \beta_P|A$ with radius outside γ' . But the radii of all balls in $b \in \beta_P|A$ are anyhow outside $\bar{\gamma}'$.

This shows that we can find d defined over A' . But d is coded, and the code is in A' , thus it is automatically A , proving the claim.

Finally, since germs on closed balls are strong ([HHM]), one can find a function g_b on b with germ $d(b)$. All the hypotheses of Lemma 5.4 are satisfied, so there exists an A -definable function g with the same P -germ as r . This finishes the proof of (v).

Proof of Theorem 2.2 By Proposition 2.4 (and Porism 2.8), we have EI to the sorts $\mathbb{Q}_p, \Gamma, S_n(\mathbb{Q}_p), T_n(\mathbb{Q}_p)$. But the sorts T_n are not actually needed; if e is a lattice, then pe is itself an R' -lattice, and a coset h of pe — a typical element of $T_n(\mathbb{Q}_p)$ — can be coded by the R' -lattice in K^{n+1} generated by h and by $(0_{K^n}, 1)$.

We finish the section with some supplementary remarks.

Remark 3.7 Let $A = ACL(A) \subseteq M \models \mathbb{Q}_p$. Then every type over A extends to an $Aut(M/A)$ -invariant type.

Proof Let $A = acl(A) \subseteq M \models T$. Let $c \in M$; then $c = f(a_1, \dots, a_n)$, where $a_i \in \text{dom}(M)$, and f is 0-definable. It suffices to extend $tp(a_1, \dots, a_n/A)$ to an $Aut(M/A)$ -invariant type. If $tp(c/M)$ and $tp(d/Mc)$ are $Aut(M/A)$ -invariant, then so is $tp(cd/M)$; so it suffices to show that $tp(a_i/A_i)$ extends to an $Aut(M/A_i)$ -invariant type for each i , where $A_i = DCL(Aa_1 \dots a_{i-1})$. But (cf. the proofs of Proposition 2.4(i) and (ii)) we have inductively that $A_i = ACL(A_i)$, so Corollary 4.7 applies.

Remark 3.8 Rigidity of finite sets fails for the theory of the p -adics (in the sorts S_n , or just in the ball sorts.)

Proof As the value group is stably embedded, one can find a non-identity automorphism σ fixing the value group. So $|\sigma(x)| = |x|$, and also $x, \sigma(x)$ have the same angular coefficient. Take a with $val(a) > 0$, with $\sigma(a) \neq a$. Let $\gamma = |\sigma(a) - a| < |a|$, $a.c.(\sigma(x) - x) = \alpha$. Then $|\sigma^2(a) - \sigma(a)| = \gamma$, $a.c.(\sigma^2(a) - \sigma(a)) = \alpha$, etc. As $p \cdot \alpha = 0$ in the residue field, $(\sigma^p(a) - a) = \sum_{i=0, \dots, p-1} (\sigma^{i+1}(a) - \sigma^i(a))$ has “absolute value $\leq \gamma$ and γ th coefficient $p\alpha$,” i.e. it has absolute value $\delta < \gamma$. Thus in the ring $\mathcal{O}/\delta$, the image of a is not a fixed point, but has an orbit of size p under σ . This set of size p is not rigid.

4 Extendible 1-types in non-principal balls

Let K be a valued field, with maximal ideal $\mathcal{M}$. We assume given also a system $\mathcal{S}$ of homomorphisms $r : K^* \rightarrow K_r$, such that each $r \in \mathcal{S}$ vanishes on $1 + \mathcal{M}^\nu$ for some $\nu = \nu(r) \in \mathbb{N}$. We assume that $(K, +, \cdot, ^{-1}, r : K \rightarrow K_r, \dots)_{r \in \mathcal{S}}$ admits elimination of quantifiers; the $\dots$ refer to additional relations on the sorts K_r , and possibly additional sorts. Assume Γ is embedded and stably embedded. Let

$$Q_R = \{(u, v, v', \dots) : (\exists x \in K) |x| = u, r(x) = v, r'(x) = v', \dots\}_{r \in \mathcal{S}}.$$

For definiteness, we assume that the value group has a convex subgroup isomorphic to $\mathbb{Z}$. Let λ_0 denote an element of maximal absolute value < 1 . Write $|x| \ll |y|$ if $|x| < |y\lambda_0^m|$ for all $m = 1, 2, \dots$. If neither $|x| \ll |y|$ nor $|y| \ll |x|$, write $x \sim y$.

We further assume that every definable subset of Q_R is coded, and moreover that germs of subsets of Q_R are strong, i.e: let $U \subseteq Q_R$. Let $\gamma_1 > \gamma_2 > \dots$ be in $\Gamma(A)$. Let $P_\Gamma = \{\delta \in \Gamma : (\forall n) \delta < \gamma_n\}$. Assume that for every conjugate U' of U over A , for some n , and some $\delta \in P_\Gamma(A)$, $\delta < x < \gamma_n \Rightarrow (x \in U \equiv x \in U')$. Then there exists U'' defined over A , $\delta \in P_\Gamma(A)$, n such that $\delta < x < \gamma_n \Rightarrow (x \in U \equiv x \in U'')$.

For the p -adics, this holds, according to Macintyre, if we take the valuation map together with the maps K^*/K^{*m} into appropriate finite groups K_r . In these finite groups, every element is named, so Q_R is just a disjoint union of copies of Γ . (This description is over-elaborate for the p -adics, and is given with a view to their ultraproducts.)

Observe that $|x - y| << |x - z|$ implies $r(x - z) = r(y - z)$ ($r \in \mathcal{S}$). Indeed $(y - z)/(x - z) = 1 + (y - x)/(x - z) \in 1 + \mathcal{M}^\nu$.

Notation 4.1 If $w \in \mathbb{B}(M)$, $x \in \text{dom}(M)$, $x \notin w$, the absolute value $|x - y|$ takes the same value for all $y \in w$. We denote it $|x - w|$. By $\text{rad}(w)$ we denote the supremum of $|a - b|$, $a, b \in w$.

If moreover $r \in \mathcal{S}$, $|x - w| |\lambda_0|^b > \text{rad}(w)$, $b > \nu(r)$, then $r(x - y) = r(x - y')$ for $y, y' \in w$. We write $r(x - w) = r(x - y)$ in this case.

Definition 4.2 Let f be a A -definable function. A partial type p over A is complete over A *relatively to f* if the natural map: $tp(c/A) \mapsto tp(f(c)/A)$ is *injective* on the set of extensions of p to $S(A)$. Similarly for several such f .

Remark 4.3 $p(x)$ is complete over A relatively to f iff for every formula $\phi(x)$ over A , there exists a formula $\theta(u)$ over A , such that $p \Rightarrow (\phi(x) \equiv \theta(f(x)))$.

For the rest of the section [check!] we consider the generic type of an ∞ -definable ball. Let $\bar{b} = \{b_n : n = 1, 2, \dots\}$ be a strictly descending sequence of balls in $\mathbb{B}(K)$. Let $P = P_{\bar{b}} = \bigcap_{n=1,2,\dots} b_n$. Let $P_\Gamma = \{\gamma \in \Gamma : (\forall n \in \mathbb{N}) \gamma < \text{rad}(b_n)\}$. For any A with $b_n \in DCL(A)$, let $q_{\bar{b}}|A$ be the conjunction of $P_b(x)$ with:

$$\{x \notin w \ \& \ |x - w| > \gamma : w \in \mathbb{B} \cap DCL(A), w \subset \bigcap_n b_n, \gamma \in P_\Gamma(A) = P_\Gamma \cap DCL(A)\}$$

Lemma 4.4 Let $A \subseteq \text{dom}(M)$. Fix $a \in \mathbb{B}(A)$ with $a \subset b_n$ for each n . Then $q_{\bar{b}}|A$ is complete relative to $|x - a|$ and to $r(x - a)$ ($r \in \mathcal{S}$).

Proof Taking into account quantifier elimination, we must show the following: let N be an elementary extension of M , let $c, c' \in N$ be two realizations of $q = q_{\bar{b}}|A$, such that $(|c - a|, r(c - a), r'(c - a), \dots)$ has the same type over A as $(|c' - a|, r(c' - a), r'(c' - a), \dots)$ ($r, r', \dots \in S$). Then the substructures $A(c), A(c')$ generated by c, c' over A are isomorphic over A .

Extend the valuation from N to N^{alg} , the algebraic closure of N . Extend each $r \in S$ to a map with kernel $Ker(r)(1 + \lambda_0^{\nu(r)} \mathcal{O}_{N^{alg}}) \subseteq N^{alg*}$.

(Note that if $a \in Ker(r), b \in \mathcal{O}_{N^{alg}}$, and $a(1 + t^{\nu(r)}b) \in N$, then $a(1 + t^{\nu(r)}b) \in Ker(r)$: indeed either $a = 0$ or $1 + t^{\nu(r)}b \in N$, and in the latter case, $b \in N$, and $|b| \leq 1$ so $b \in \mathcal{O}$, and thus $(1 + t^{\nu(r)}b) \in Ker(r)$.)

So we still have: $|x - y| \ll |x - z|$ implies $r(x - z) = r(y - z)$.

Then it suffices to show that $A^{alg}(c), A^{alg}(c')$ are A^{alg} -isomorphic, by an isomorphism commuting with the extensions of the maps r . (One can then restrict the isomorphism to $M(c)$.)

Let $\gamma = |a - c|, \gamma' = |a - c'|$. Then γ, γ' realize the same type in the value group over A ; so we may assume $\gamma = \gamma'$. Similarly we may assume $r(a - c) = r(a - c')$ for each $r \in S$ (the corresponding $*$ -types agree.)

Take any $b \in A^{alg}$. If $b \notin b_n$, for some n , then $|c - b| = |a - b| = |c' - b|$. Moreover, $|c - c'| < rad(b_{n+k}) \leq rad(b_n)|\lambda_0|^k < |c - b||\lambda_0|^k$; and it follows that $r(c - b) = r(c' - b)$ for $r \in S$.

If $b \in b_n$ for each n , then $|a - b| \ll rad(b_n)$ for each n . Note that $m|a - b| = |e|$ for some $e \in A$, so $|a - b| \in DCL(A)$. Thus $|a - c| \gg |a - b|$. Thus $r(c, b) = r(a, b)$. We have $|a - c'| = |a - c|$, so $|b - c'| = |a - c'| = |a - c| = |b - c|$, and $r(c', b) = r(a, b) = r(c, b)$. So in either case $|c - b| = |c' - b|$, $r(b - c) = r(b - c')$ ($r \in S$).

As any rational function g over A is a ratio of products of constant or linear polynomials, it follows that $|g(c)| = |g(c')|, r(g(c)) = r(g(c'))$ in general. This proves the lemma.

Lemma 4.5 *Let $A \subseteq M$, $A = DCL(A)$. Then for any formula $\phi(x)$ over A , either $q_{\bar{b}}|A \Rightarrow \phi$ or $q_{\bar{b}}|A \Rightarrow \neg\phi$ or there exists a formula $\theta(u, v, v', \dots)$ over A , and proper sub-balls $e \subset e' \subset P$ defined over A , such that*

$$x \notin e' \Rightarrow (\phi(x) \equiv \theta(|x - e|, r(x - e), \dots))$$

Thus:

(a) *Let $a \in \mathbb{B}(A)$ with $a \subset b_n$ for each n . Then $q_{\bar{b}}|A$ is complete relative to $|x - a|$ and to $r(x - a)$ ($r \in S$).*

(b) *If there is no a as in (a), then $q_{\bar{b}}|A$ is complete.*

Proof Let $\phi(x)$ be formula over A . Let $\bar{u} = (u, v, v', \dots)$ be a sequence of variables appropriate for $|x|, r(x), r'(x), \dots$. Applying Lemma 4.4 over M (enlarged if necessary so that $P(M)$ is nonempty), for any $a \in P(M)$, there exists a formula $\theta = \theta_a$ such that

$$q_{\bar{b}}|M \Rightarrow (\phi(x) \equiv \theta(|x - a|, r(x - a), \dots))$$

Note that two conjugates of $\theta(u, v, v', \dots)$ over A must agree, on elements $(u, v, v', \dots) \in Q_R$ such that u stays away from some proper sub-segment of P_Γ . (Say θ, θ' are defined over A' ; if $u > \delta$ for any $\delta \in P_\Gamma(A')$, pick x with $|x - a| = u, r(x - a) = v, \dots$; then $|x - a'| = u, r(x - a') = v, \dots$, and so $\theta(u, v, v', \dots)$ iff $\phi(x)$ iff $\theta'(u, v, v', \dots)$.) By the strong coding property assumed for Q_R , θ can be taken to be defined over A .

Now a finite union of proper sub-balls of P is contained in a single proper sub-ball. So there exists a proper sub-ball e of P such that $x \notin e \Rightarrow (\phi(x) \equiv \theta(|x - a|, r(x - a), \dots))$. Let e' be the ball containing e and finitely many sizes bigger; but big enough that for each $r \in \mathcal{S}$ mentioned in θ , if $y \notin e'$ then $r(y, e)$ is well-defined. Then:

$$x \in P, x \notin e' \Rightarrow (\phi(x) \equiv \theta(|x - e|, r(x - e), \dots)). \quad (\#)$$

Let $\alpha = \text{rad}(e)$. So $\text{rad}(e') = \alpha|\lambda_0|^{-l}$ for some integer l ; we fix l , so that e' is a function of e .

If $q_{\bar{b}}|A \Rightarrow \neg\phi$ or $q_{\bar{b}}|A \Rightarrow \phi$, we are done. Otherwise, there are $a_0, a_1 \models q_{\bar{b}}|A$ with $\phi(a_1), \neg\phi(a_0)$. Let f be the smallest ball containing a_0, a_1 . Let f' be the ball containing f of radius l notches more than that of f . Then there cannot be (e, e') satisfying $(\#)$ with f', e' disjoint. Otherwise, $|a_0 - e| = |a_1 - e|$ and $r(a_0 - e) = r(a_1 - e)$ for any r mentioned in θ ; so $\theta(|a_0 - e|, r(a_1 - e), \dots)$ holds iff $\theta(|a_1 - e|, r(a_1 - e), \dots)$, and thus $\phi(a_0)$ iff $\phi(a_1)$, a contradiction.

But any conjugate of (e, e') over A has $(\#)$. Thus, no two conjugates of e' are disjoint. Hence they form a linearly ordered chain; an element of such a chain is determined by its radius. So we can write $e' = e'(\beta)$; $e'(\beta)$ is a ball of radius β , and e' is an A -definable function of β . Let $e''(\beta)$ be l notches bigger; then $(\#)$ holds of $(e'(\beta), e''(\beta))$.

The domain of these functions can be taken to be an A -definable interval in Γ ; such that for each β' in this interval, $(\#)$ holds of $e'(\beta'), e''(\beta')$. Take the least β_0 in the interval (here we use the embeddedness and stable embeddedness of Γ ; nonempty definable subsets of Γ have least elements. We could say instead: some model of the theory has well-ordered positive Γ .) Now

$e'(\beta_0) \subset e'(\beta) \subset P$, so $e'(\beta_0) \subset P$, and thus $e''(\beta_0) \subset P$. As β_0 is defined over A , we have found the two balls as required.

Lemma 4.6 *Assume Γ is definably well ordered (every nonempty definable subset has a least element). Let $A \subseteq M, \bar{b} \in DCL(A)$ a properly decreasing sequence of balls. Let q_r be a complete type over A extending Q_R . Assume q_r implies $u < \text{rad}(b_n)$ for each n , and $\gamma < u$ for any $\gamma \in DCL(A)$, $\gamma < \text{rad}(b_n)$ (each n .) Then*

$$q_{\bar{b}} \cup \bigcup_{a \in P_{\bar{b}}(M)} q_r(|x - a|, r(x - a), r'(x - a), \dots)$$

is consistent.

Proof We may assume M has an element a' with $a' \in b_n$ for each n . Let $G = \{\gamma \in M, (\forall n)(\gamma < \text{rad}(b_n))\}$. Note that q_r is consistent with $\{\gamma < u : \gamma \in G\}$.

(For any $\gamma \in G$, if $q_r \models u \leq \gamma$, then some $\psi \in q_r$ is bounded below γ ; but then the least upper bound γ' of $\{u : \psi < u\}$ is in $DCL(A)$, so $q_r \models \gamma' < u$, a contradiction.)

Let c' be such that $(|c'|, r(c'), \dots) \models q_r \cup \{\gamma < u : \gamma \in G\}$.

Let $d = a' + c'$. Clearly $q_{\bar{b}}(d)$. Moreover for any $a \in P_{\bar{b}}(M)$, $|d - a'| = |c'| \gg |a - a'|$. Thus $|d - a| = |d - a'|$ and so, as observed before, $r(a' - d) = r(a - d) = r(c')$. Thus d realizes the type in question.

Corollary 4.7 *Let $M \models Th(\mathbb{Q}_p)$, $A \subseteq M$, $ACL(A) \cap \mathbb{B} \subseteq A$. Let $c \in \text{dom}(M)$. Then $tp(c/A)$ extends to an $\text{Aut}(M/A)$ -invariant type.*

Proof Let $W(c; A) = \{w \in \mathbb{B}(A) : c \in w\}$, $P(c; A) = \bigcap W(c; A)$. As the residue field of M is finite, $P(c; A)$ cannot reduce to a single ball (that ball would be the union of finitely many proper sub-balls, each in $ACL(A) \cap \mathbb{B}$, hence in A .) Thus Lemma 4.5 applies to $P = P(c; A)$. Let q_P be the relatively complete type described there, for P . Let q_r be any $\text{Aut}(M/A)$ -invariant type, extending Q_R and $u < \text{radius}(b)$ ($b \in W(c; A)$), $\alpha < u$ ($u \in P_\Gamma$). By Lemma 4.6, $q^* = q_P(x) \cup \bigcup_{a \in P} q_r(|x - a|, r(x, a), \dots)$ is consistent. Clearly q^* is $\text{Aut}(M/A)$ -invariant. The facts that it is complete, and that it extends $tp(c/A)$, both follow from Lemma 4.5.

Let N_n be the group of matrices of the form $I_n + b$, where I_n is the identity matrix in GL_n , and b is matrix in $B_n(\mathcal{O})$ with all entries having absolute value $<< 1$.

Thus $N_n(\mathbb{Q}_p) = B_n(\mathbb{Q}_p) \cap \cap_m (I_n + p^m M_n(\mathcal{O}))$. This can be viewed as the identity component of $B_n(\mathbb{Z}_p)$.

Lemma 4.8 (N_n has a right-generic type.) *There exists an $\text{Aut}(M)$ -invariant type $p|M$ of matrices $a \in N_n$, also invariant under right multiplication: if $b \in A \subseteq N_n$, and $c \models p|A$, then $cb \models p|A$. p is complete relative to the norms and r_n -values of the entries. (Or, if Q_R has an $\text{Aut}(M)$ -invariant 1-type, p can be taken to be complete.)*

Proof Let $q = q_{\bar{b}}$ be the 1-type over M given as in Lemma 4.5, with respect to the cut: $\bar{b} = (|\lambda_0|, |\lambda_0^2|, \dots)$. (Or, let q be any complete, $\text{Aut}(M)$ -invariant extension of this type.)

Let p be the type of upper-triangular matrices obtained by iterating q $n(n+1)/2$ times, using the lexicographic order on the matrix entries, and adding 1 on the diagonal: thus if $(a) \in M_n$, then $I_n + (a) \models p|A$ iff $a_{11} \models q|A$, $a_{12} \models q|DCL(A, a_{11})$, $\dots$, $a_{22} \models q|DCL(A, a_{11}, \dots, a_{1,n})$, $\dots$, $a_{n,n} \models q|DCL(A, a_{11}, \dots, a_{n,n-1})$, while $a_{ij} = 0$ for $i > j$.

The fact that p is an $\text{Aut}(M)$ -invariant type of elements of N_n is clear. As for the right translation invariance, let $I_n + (b) \in N_n$; we have to show that $(I_n + (a))(I_n + (b)) = I_n + (a) + (b) + (a)(b) \models p$. Let $(d) = (a) + (b) + (a)(b)$. Then $d_{11} = a_{11} + b_{11} + a_{11}b_{11}$. We have

$$\mathbb{Z} < \text{val}(a_{11}) << \text{val}(b_{11}) < \text{val}(a_{11}b_{11}).$$

So $\text{val}(d_{11}) = \text{val}(a_{11})$, $r(d_{11}) = r(a_{11}) = 1$ (for $r \in \mathcal{S}$); it follows that $d_{11} \models q$. Similarly

$$d_{12} = a_{12} + b_{12} + a_{11}b_{12} + a_{12}b_{22};$$

here a_{12} has strictly smaller value than any of the other summands, so again $\text{val}(d_{12}) = \text{val}(a_{12})$. Thus $d_{12} \models q|DCL(A, a_{11})$. But since $b \in DCL(A)$, $d_{11} \in DCL(A, a_{11})$, so $d_{12} \models q|DCL(A, d_{11})$. Continuing in this way we see that $(d) \models p$.

Corollary 4.9 *Let R be a (left or right) coset of N_n in $B_n(K)$. Then there exists an $\text{Aut}(M/R)$ -invariant type of elements of R .*

Proof Say R is a right coset. Pick $g \in R$, let p be the right- N_n -invariant type of Lemma 4.8, and let ${}^g p = tp(gc/A)$, where $c \models p|DCL(Ag)$. Then ${}^g p = {}^{gh} p$ for $h \in N_n$, since p is N_n -invariant. Thus any automorphism fixing R must fix ${}^g p$.

Corollary 4.10 *Let $M \models Th(\mathbb{Q}_p)$, $e \in S_n(M)$, $E = ACL(e) \cap M$. Then there exists an $Aut(M/E)$ -invariant type of bases for e .*

Proof It was noted above [ref?] that any lattice e has a triangular basis; this basis can be viewed as the set of columns of a matrix in $B_n(K)$. Let b, b' be two such bases, and suppose $b' = \sigma(b)$, $\sigma \in Aut(M/E)$. Then as $e/p^m e$ is finite for all m , the columns of b, b' must be in the same coset of $p^m e$ for each m . Thus if we write $b' = b(a)$ with $(a) \in B_n(\mathcal{O})$, then $(a) = I_n$ modulo p^m for each m , so $(a) \in N_n$. Thus $Aut(M/E)$ preserves the coset bN_n . So it suffices to take the $Aut(M/bN_n)$ -invariant type of elements of bN_n guaranteed by Corollary 4.9.

5 Algebraically Closed Valued Fields

Here we recall some facts about algebraically closed valued fields, essentially from [HHM].

Let K be a (sufficiently saturated) algebraically closed valued field, $A = acl(A) \subset K$. Let P be a ball, or an infinite intersection of balls, defined over A . If $P = \bigcap P_n$, then the cut in Γ below $\{rad(P_n) : n\}$ is referred to as $rad(P)$.

There is an associated $Aut(K/A)$ -invariant 1-type α_P of elements of P : $c \in \alpha_P|B$ if $c \in P$, and there is no B -definable proper sub-ball b of P with $c \in b$. We say that c is generic in P over B if $c \in \alpha_P|B$.

When P is open or strictly ∞ -definable, there is also an associated $Aut(K/A)$ -invariant type of “large, centered” closed balls in P ; we denote it β_P , and refer to a generic closed ball of P . Namely, for $A \subseteq B = acl(B)$, $b \in \beta_P|B$ if b is a closed ball containing any B -definable proper sub-ball of P , and of radius bigger than any $\gamma \in \Gamma(B)$ with $\gamma < rad(P)$.

It follows that b is contained in no proper B -definable sub-ball of P ; so a B -generic $a \in b$ is also B -generic as an element of P .

Thus $tp(b/B)$ can also be described as follows: first pick a generic $c \in P$; then pick a generic $\alpha < rad(P)$, and let b be the closed ball around c of radius $|\alpha|$.

If P has an A -definable proper sub-ball b_0 , then there is no need for c : b is the ball around b_0 with radius $|\alpha|$. In this case $dcl(b, B) = dcl(\alpha, B)$.

Definition 5.1 Let r be a function (defined over $A' = acl(A') \supseteq A$), with $r(x)$ a finite set. Let q be an $Aut(K/A)$ -invariant type. r is *irreducible* on q

if for generic $a \in q$ (over A'), all elements of $r(a)$ are $\text{Aut}(K/A'a)$ -conjugate. Equivalently, there is no r' defined over A' such that for generic $c \in q$, $r'(c)$ is a nonempty proper subset of $r(c)$.

Lemma 5.2 *The notion of irreducibility on β_P does not depend on A' ; increasing A' will not change it. (For sequentially generic types, this is the Stationarity Lemma of HHM-2.)*

Proof Let r be a function defined over $A' = \text{acl}(A') \supseteq A$, and assume that for $c \in \alpha_P|A$, $\text{Aut}(K/A'c)$ acts transitively on $r(c)$. We will show the same is true for any $B \supseteq A'$.

If P has a proper sub-ball defined over A' , then $\beta_P|A'$ is definably equivalent to the type of a generic $\alpha < \text{rad}(P)$, so [the Stationarity Lemma of HHM-2] applies directly. If P has no sub-ball, then $\Gamma(A') = \Gamma(A'c)$, if $c \in P$ is generic over A' . Let $s(x, u) = r(b(x, u))$, where $b(x, u)$ is the ball around x of radius u . Let $b \in \beta_P(A')$, pick a generic point c of b (over $A'b$), and let α be the radius of b . Then $\text{tp}(c/A'b)$ extends to an $\text{Aut}(K/A'b)$ -invariant type, so in particular every $A'b$ -elementary permutation of $r(c)$ is also elementary over $A'bc$, hence over $A'c\alpha$; thus s is irreducible (over A'). Let q be the sequentially generic type of pairs (x, u) , with $x \in P$ generic, and then $u < \text{rad}(P)$ generic over x . Then by [the Stationarity Lemma of HHM-2], for any B , if $(c', \alpha') \models q|b$, $b' = b(c', \alpha')$, then $s(c', \alpha') = r(b')$ is transitive under $\text{Aut}(K/Bc'\alpha')$, hence certainly under $\text{Aut}(K/Bb')$. As $b' \in \beta_P|B$, this proves the lemma.

Lemma 5.3 *Let r be irreducible on P . Then for a generic closed ball $C \subset P$, $r|C$ is irreducible on C .*

Proof Let B be a base set with $A \subseteq B = \text{acl}(B)$, r defined over B , and such that P has a point $b_0 \in B$. Let $C \in \beta_P|B$. Then $b_0 \in C$, and C is the ball around b_0 of some radius α ; C is defined over B, α . Let $c \in C$ be a generic point (over B, α). Then c is a generic point of P over B . Thus the elements of $r(c)$ are $\text{Aut}(K/B, c)$ -conjugate. But $\alpha \in \text{dcl}(B, c)$ since $|b_0 - c| = \alpha$. So the elements of $r(c)$ are $\text{Aut}(K/B, \alpha, c)$ -conjugate. Let $B' = \text{dcl}(B, \alpha) = \text{dcl}(B, \langle C \rangle)$. Then by HHM, $\text{acl}(B') = B'$. $\text{Aut}(K/B', c)$ is transitive on $r(c)$, so proving that $r|C$ is irreducible on C .

Lemma 5.4 *Assume that for $C \in \beta_P|A$, there is given a function g_C on C , with g_C defined over $A, \langle C \rangle$. Assume moreover that there exists a K -definable function r such that for $C \in \beta_P(A)$, $r|C, g_C$ have the same C -germ.*

Then there exists an A -definable function g on P , such that for a generic closed ball C of P , $g|C, g_C|C$ have the same C -germ. (Moreover, r, g have the same P -germ.)

Proof First suppose P has a proper sub-ball c_0 defined over A . In this case the generic closed balls C have the form $C(\alpha)$, where α is generic in an appropriate cut in Γ , and $C(\alpha)$ is the closed ball around c_0 of radius α . Let $C'(\alpha)$ be the open ball around c_0 of radius α , and let $A(\alpha) = C(\alpha) \setminus C'(\alpha)$. Then the sets $A(\alpha)$ are disjoint, and we can form a function g with $g(t) = g_C(t)$ whenever $t \in A(\alpha)$, $C = C(\alpha)$ (gluing on annuli.) It is then clear that $g|C, g_C|C$ have the same C -germ.

The function r was not needed so far; but if it is given, then r, g can at worst differ on a proper-sub-ball of P (otherwise they agree on at most a proper sub-ball; for a generic closed C containing that sub-ball properly, $r|C, g|C$ will not have the same C -germ, contradiction.)

If on the other hand P has no A -definable proper sub-balls, P forms a complete type over A ; and the type over A of a nested tuple of sub-balls of P , is determined by the type of their radii.

Let $C_0 \in \beta_P(A)$, $C \in \beta_P(A, \langle C_0 \rangle)$. r must agree with g_C away from some finite union F of proper sub-balls of C . Find $C' \subseteq C$, $C' \cap F = \emptyset$, $\text{rad}(C') = \text{rad}(C_0)$; then $C \in \beta_P(A, \langle C' \rangle)$. As $r|C' = g_C|C'$, we have: $g_{C'}, g_C|C'$ agree generically. But C' has no proper balls defined over $\text{acl}(A, \langle C' \rangle, \langle C \rangle)$ (since there are infinitely many other sub-balls with the same radii, and by the remark on types of nested balls [label?].) Thus $g_{C'}, g_C|C'$ agree with no exceptions.

Now $C' \in \beta_P(A)$, $C \in \beta_P(A, \langle C' \rangle)$. Whenever $C_1 \subset C_2 \in \beta_P(A)$, consider a $C^* \in \beta_P(A, \langle C_1 \rangle, \langle C_2 \rangle)$; then $g_{C^*}|C_i = g_{C_i}$, so $g_{C_2}|C_1 = g_{C_1}$. Thus $g = \cup_{C \in \beta_P(A)} g_C$ is a function, satisfying the requirements.

Finally we repeat verbatim a lemma and proof from [HHM-2].

Lemma 5.5 *Let $(P, Q; R)$ be a structure; $R \subseteq P^2 \times Q$. Assume:*

- (i) *For any $a \in P$, $\text{acl}(a) \cap Q = \emptyset$*
- (ii) *For any $a \neq b \in P$, $R(a, b) = \{c \in Q : R(a, b, c)\}$ is a finite subset of Q , of bounded size. Assume for simplicity $R(a, b) = R(b, a)$.*
- (iii) *For distinct $a, b, c \in P$, $R(a, b) \subseteq R(a, c) \cup R(b, c)$.*

Then $R(a, b) = \emptyset$.

Proof We may assume the structure is $\aleph_0$ -homogeneous. Pick $a, b \in P$ with $R(a, b)$ of maximal size n . Let $F = R(a, b)$.

Claim 5.6 For any $b' \in P$, if $b' \neq b$ and $R(b, b') \neq \emptyset$ then $R(b, b') \cap F \neq \emptyset$.

Proof For $b' = a$, this is clear (unless $n = 0$ and we are done.) Let $b' \neq a$. Suppose $R(b, b') \cap R(a, b) = \emptyset$. Then $R(b, b') \subseteq R(a, b')$ and $R(a, b) \subseteq R(a, b')$, by (ii). But by maximality of n , this forces $R(a, b) = R(a, b')$, so $R(b, b') = \emptyset$.

Now $F \cap \text{acl}(b) = \emptyset$. By Neumann's Lemma, there exist disjoint conjugates $F_1, \dots, F_{n+1}$ of F over b . So Claim 5.6 as stated applies to any F_i in place of F . Pick any $b' \in P$, $b' \neq b$, with $R(b, b') \neq \emptyset$ (e.g. $b' = a$); then by Claim 5.6, $R(b, b') \cap F_i \neq \emptyset$, for each i . But the $n + 1$ sets F_i are disjoint; so $R(b, b')$ has at least $n + 1$ points, a contradiction.

6 Rationality

We will need the following consequence of EI: for any 0-definable set $\mathcal{D}$ and any 0-definable equivalence relation $\mathcal{E}$ on $\mathcal{D}$, there is a 0-definable map f from $\mathcal{D}$ to a product of sorts $M_1 \times \dots \times M_n$ for some n such that for every $x, y \in \mathcal{D}$, we have $x\mathcal{E}y$ if and only if $f(x) = f(y)$ (see [HHM], Section 1). It is not hard to show that if a theory admits EI then it still does so after adding some constants to the language, so we may replace all of the occurrences of “0-definable” by “definable” in the previous sentence.

Lemma 6.1 *Let $N \in \mathbb{N}$ and let μ be the additive Haar measure on $\mathbb{Q}_p^N$, normalised so that $\mu(\mathbb{Z}_p^N) = 1$. For any $A \in \text{GL}_N(\mathbb{Q}_p)$, we have*

$$\mu(A \cdot \text{GL}_N(\mathbb{Z}_p)) = |\det A|^N. \quad (2)$$

Proof Since $\mathbb{Z}_p^{N^2} - \text{GL}_N(\mathbb{Z}_p)$ has Haar measure zero (it is the vanishing set of a nontrivial polynomial), we may replace $\text{GL}_N(\mathbb{Z}_p)$ with $\mathbb{Z}_p^{N^2}$ in Eqn. (2). We identify $\mathbb{Z}_p^{N^2}$ with the free $\mathbb{Z}_p$ -module D on the matrices E_{ij} for $1 \leq i, j \leq N$, where the (i, j) -entry of E_{ij} is one and the other entries are zero.

Let $A \in \text{GL}_N(\mathbb{Q}_p)$. Let A_1 be a matrix obtained either by exchanging two columns of A , or by adding some $\mathbb{Z}_p$ -multiple of one column of A to

another column of A . Then $|\det A| = |\det A_1|$, and we have $A.D = A_1.D$, so $\mu(A.D) = \mu(A_1.D)$. Now fix $k \in 1, \dots, N$ and let A_2 be the matrix obtained from A by multiplying the k th column of A by some nonzero $\lambda \in \mathbb{Q}_p$. Then $\det A_2 = \lambda \det A$. Suppose that $|\lambda| \geq 1$. Let $D_k(\lambda)$ be the free $\mathbb{Z}_p$ -module generated by the E_{ij} for $1 \leq i, j \leq N$ and $i \neq k$, together with the λE_{kj} for $1 \leq j \leq N$. We have $A_2.D = A.D_k(\lambda)$. Now $|A.D_k(\lambda)/A.D| = |D_k(\lambda)/D| = |\lambda|^N$, so $A.D_k(\lambda)$ is the union of $|\lambda|^N$ disjoint translates of $A.D$. It follows that $\mu(A_2.D) = |\lambda|^N \mu(A.D)$. The same formula holds by a similar argument if $|\lambda| < 1$. Since any element of $GL_N(\mathbb{Q}_p)$ can be obtained from the identity matrix by elementary column operations, Eqn. (2) follows.

By a *definable family* $R = (R_l)_{l \in \mathbb{Z}^r}$ of subsets of $\mathbb{Q}_p^N$ we mean a definable subset R of $\mathbb{Q}_p^N \times \mathbb{Z}^r$, and we write R_l for the fibre above l of the projection from R to $\mathbb{Z}^r$. By a *definable family* $E = (E_l)_{l \in \mathbb{Z}^r}$ of equivalence relations on R we mean a definable equivalence relation E on R such that for every $x, y \in R$, if xEy then there exists $l \in \mathbb{Z}^r$ such that $x, y \in R_l$. We then have a definable equivalence relation E_l on R_l for every l , and by a slight abuse of notation we can regard $(E_l)_{l \in \mathbb{Z}^r}$ as a definable family of subsets of $\mathbb{Q}_p^{2N}$. Let $\mathbb{N}_0$ denote the set of nonnegative integers. Then $\mathbb{N}_0^r$ is a definable subset of $\mathbb{Z}^r$, so it makes sense to talk of definable families $R = (R_l)_{l \in \mathbb{N}_0^r}$, etc.

Now we come to the main result of this section. We extend the definition of $\mathbb{Q}$ -rationality to power series in several variables in the obvious way.

Theorem 6.2 *Let $R = (R_l)_{l \in \mathbb{N}_0^r}$ be a definable family of subsets of $\mathbb{Q}_p^N$. Let $E = (E_l)_{l \in \mathbb{N}_0^r}$ be a definable family of equivalence relations on $(R_l)_{l \in \mathbb{N}_0^r}$. Suppose that for each $l \in \mathbb{N}_0^r$, the set of equivalence classes R_l/E_l is finite. Let $a_l = |R_l/E_l|$. Then the power series*

$$\sum_{l \in \mathbb{N}_0^r} a_l t^l \in \mathbb{Q}[[t_1, \dots, t_r]]$$

is $\mathbb{Q}$ -rational.

Proof By EI, there exists $m \in \mathbb{N}$, $M_1, \dots, M_s \in \mathbb{N}$ for some $s \in \mathbb{N}$, and a definable function $f': R \rightarrow \mathbb{Q}_p^m \times \mathcal{S}_{M_1} \times \dots \times \mathcal{S}_{M_s}$ such that for every $x, y \in R$, $xEy \iff f'(x) = f'(y)$. Define $f: R \rightarrow \mathbb{Q}_p^m \times \mathcal{S}_{M_1} \times \dots \times \mathcal{S}_{M_s} \times \mathbb{N}_0^r$ by $f(x) = (f'(x), l)$ if $x \in R_l$.

The determinant function on $GL_M(\mathbb{Q}_p)$ descends to give a well-defined definable function $\det: \mathcal{S}_M \rightarrow \mathbb{Q}_p$. Define $\mathcal{S}_M^+ = \{x \in \mathcal{S}_M \mid \log_p |\det x| \geq 0\}$

and $\mathcal{S}_M^- = \{x \in \mathcal{S}_M \mid \log_p |\det x| < 0\}$. The map $B \mapsto (B^T)^{-1}$ (where B^T denotes the transpose of B) descends to give a well-defined definable bijection $\alpha_M: \mathcal{S}_M \rightarrow \mathcal{S}_M$, mapping $\mathcal{S}_M^-$ into $\mathcal{S}_M^+$. The preimages $f^{-1}(\mathbb{Q}_p^m \times \mathcal{S}_{M_1}^{\epsilon_1} \times \cdots \times \mathcal{S}_{M_s}^{\epsilon_s} \times \mathbb{N}_0^r)$, where each ϵ_i is one of the symbols $+$ or $-$, partition R into a finite disjoint union of definable pieces. On each piece we can replace f with $(\text{id}_{\mathbb{Q}_p} \times g_1 \times g_s \times \text{id}_{\mathbb{Z}^r}) \circ f$, where each g_i is $\text{id}_{\mathcal{S}_{M_i}}$ if $\epsilon_i = +$ and α_{M_i} if $\epsilon_i = -$. Thus we can assume without loss that $f(R) \subseteq \mathbb{Q}_p^m \times \mathcal{S}_{M_1}^+ \times \cdots \times \mathcal{S}_{M_s}^+ \times \mathbb{N}_0^r$.

Denote by π_M the canonical projection from $\text{GL}_M(\mathbb{Q}_p)$ to $\mathcal{S}_M$. Let $A = (A_l)_{l \in \mathbb{N}^r}$ be the definable family of subsets of $\mathbb{Q}_p^m \times \mathbb{Q}_p^{M_1^2} \times \cdots \times \mathbb{Q}_p^{M_s^2}$ consisting of tuples $(x, B_1, \dots, B_s, l) \in \mathbb{Q}_p^m \times \mathbb{Q}_p^{M_1^2} \times \cdots \times \mathbb{Q}_p^{M_s^2} \times \mathbb{N}_0^r$ such that

$$(x, \pi_{M_1}(B_1), \dots, \pi_{M_s}(B_s), l) \in f(R_l).$$

Each A_l is the union of a_l subsets of the form

$$D = \{x\} \times B_1 \cdot \text{GL}_{M_1}(\mathbb{Z}_p) \times \cdots \times B_s \cdot \text{GL}_{M_s}(\mathbb{Z}_p) \times \{l\},$$

where each $B_i \in \text{GL}_{M_i}(\mathbb{Q}_p)$.

Let $\pi: A \rightarrow \mathbb{Q}_p^{M_1^2} \times \cdots \times \mathbb{Q}_p^{M_s^2} \times \mathbb{N}_0^r$ be the projection that omits the first factor $\mathbb{Q}_p^m$, and let $A' = (A'_l)_{l \in \mathbb{N}_0^r}$ be the image of A , a definable family of subsets of $\mathbb{Q}_p^{M_1^2} \times \cdots \times \mathbb{Q}_p^{M_s^2}$. By slight abuse of notation we regard the A'_l as definable subsets of $\mathbb{Q}_p^{M_1^2} \times \cdots \times \mathbb{Q}_p^{M_s^2}$. Each A'_l is the union of a finite number a'_l of subsets D' of the form

$$D' = B_1 \cdot \text{GL}_{M_1}(\mathbb{Z}_p) \times \cdots \times B_s \cdot \text{GL}_{M_s}(\mathbb{Z}_p) \quad (3)$$

and the fibres of π are finite. As the theory of $\mathbb{Q}_p$ is algebraically bounded [ref], the fibre size is bounded by some positive integer N . For each j between 1 and N , define $A'(j) = (A'_l(j))_{l \in \mathbb{N}_0^r}$ to be the definable set $\{x \in A' \mid |\pi^{-1}(x)| = j\}$. Each $A'_l(j)$ is the disjoint union of some finite number $a'_l(j)$ of subsets of the form (3). We have $a_l = \sum_{j=1}^N a'_l(j)$ for each l , so it suffices to show that for each j , $\sum_{l \in \mathbb{N}_0^r} a'_l(j) t^l$ is $\mathbb{Q}$ -rational.

So fix j . Define functions $g_i: A'(t) \rightarrow \mathbb{Z}$ for $1 \leq i \leq s$ by $g_i(B_1, \dots, B_s, l) = \log_p(|\det B_i|^{M_i})$. By our assumption on $f(A)$, each g_i takes only nonnegative values. For each piece D' of the form (3) in $A'_l(j)$, Eqn. (2) gives

$$\int_{D'} p^{-(g_1 + \cdots + g_s)} d\mu = 1,$$

so we have

$$a'_l(j) = \int_{A'_l} p^{-(g_1+\dots+g_s)} d\mu.$$

Since the g_i take only nonnegative values, it now follows from [2], Theorem 1.5 and Theorem 1.6.1 that the power series

$$\sum_{l \in \mathbb{N}_0^r} a'_l(j) t^l \in \mathbb{Q}[[t_1, \dots, t_r]]$$

is $\mathbb{Q}$ -rational, as required.

The function $l \mapsto a_l$ is built from piecewise linear functions by multiplication, exponentiation and $\mathbb{Q}$ -linear combinations (see [2], Sections 1.5 and 1.6). This implies the following growth estimate on a_l : there exist $K > 0$ and $c_1, \dots, c_r \geq 0$ such that for every $l = (n_1, \dots, n_r) \in (\mathbb{Z}_{\geq 0})^r$, we have

$$a_l \leq K c_1^{n_1} \cdots c_r^{n_r}. \quad (4)$$

Below we consider definable families that arise in the following way. Let $\mathcal{D} \subset \mathbb{Q}_p^N$ be definable and let $\mathcal{E}$ be a definable equivalence relation on $\mathcal{D}$. Suppose that $f_1, \dots, f_r: \mathcal{D} \rightarrow \mathbb{Q}_p - \{0\}$ are definable functions such that for every $l \in \mathbb{Z}^r$, the subset $\{x \in \mathcal{D} \mid f_i(x) = p^{-l_i}\}$ is a union of equivalence classes. Then we can regard $\mathcal{D}$ as a definable family $(\mathcal{D}_l)_{l \in \mathbb{Z}^r}$ of subsets of $\mathbb{Q}_p^N$, identifying $\mathcal{D}$ with the set $\{(x, l) \in \mathcal{D} \times \mathbb{Z}^r \mid |f_i(x)| = l_i \text{ for all } i\}$, and we can regard $\mathcal{E}$ as a definable family of equivalence relations on $\mathcal{D}$.

7 Subgroup Growth

We write $\widehat{G}_p$ for the pro- p completion of a group G .

We give some examples of rational zeta functions from the theory of subgroup growth. If G is a finitely generated group then for any $n \in \mathbb{N}$, the number a_n of index n subgroups of G is finite. We use Theorem 6.2 to give alternative proofs of results of [4] and [dS] concerning rationality of power series constructed using the subgroup growth function when G is nilpotent. First we need to recall some facts about nilpotent pro- p groups, including the notion of a good basis for a subgroup of a torsion-free nilpotent group [4], Section 2; we will need these ideas in Section 8 as well.

Let Γ be a finitely generated nilpotent group and let $j: \Gamma \rightarrow \widehat{\Gamma}_p$ be the canonical map. Then $\widehat{\Gamma}_p$ is finitely generated as a pro- p group, so every

finite index subgroup of $\widehat{\Gamma}_p$ is open ([3], 1.17 Theorem) and has p power index ([3], 1.18 Lemma). Since Γ is finitely generated nilpotent, every p -power index subgroup is open in the pro- p topology on Γ (this follows from [ref: Hall], Lemma 2.6 and its corollaries). In particular, there is a bijection $H \mapsto \overline{j(H)}$ between index p^n subgroups of Γ and index p^n subgroups of $\widehat{\Gamma}_p$, and $\overline{j(H)} \simeq \widehat{H}_p$ (see [4], Proposition 1.2). For any $H \leq \Gamma$ of index p^n , we have $\Gamma/H \simeq \widehat{\Gamma}_p/\overline{j(H)}$.

[Emphasise that Γ is nilpotent.]

Let Δ be a finitely generated torsion-free nilpotent group. A *Mal'cev basis* is a tuple $a_1, \dots, a_R$ of elements of Δ such that any element of Δ can be written uniquely in the form $a_1^{\lambda_1} \cdots a_R^{\lambda_R}$, where the $\lambda_i \in \mathbb{Z}$. We call the λ_i *Mal'cev co-ordinates*. Moreover, group multiplication and inversion in Δ are given by polynomials in the λ_i with coefficients in $\mathbb{Q}$, as is the map $\Delta \times \mathbb{Z} \rightarrow \Delta$, $(g, \lambda) \mapsto g^\lambda$. We may regard the a_i as elements of the pro- p completion $\widehat{\Delta}_p$, and analogous statements hold, except that λ and the Mal'cev co-ordinates λ_i now belong to $\mathbb{Z}_p$. In particular, the map $j: \Delta \rightarrow \widehat{\Delta}_p$ is injective and we may identify $\widehat{\Delta}_p$ with $\mathbb{Z}_p^R$.

Now let H be a finite index subgroup of $\widehat{\Delta}_p$, of index p^n , say. A *good basis for H* is an R -tuple $h_1, \dots, h_R \in H$ such that every element of H can be written uniquely in the form $h_1^{\lambda_1} \cdots h_R^{\lambda_R}$ ($\lambda_i \in \mathbb{Z}_p$), and satisfying an extra property which does not concern us here. We say that $h_1, \dots, h_R \in \widehat{\Delta}_p$ is a *good basis* if it is a good basis for some finite index subgroup H of $\widehat{\Delta}_p$. For each i , we can write

$$h_i = a_1^{\lambda_{i1}} \cdots a_R^{\lambda_{iR}} \quad (5)$$

and we recover $|\widehat{\Delta}_p: H|$ from the formula

$$|\widehat{\Delta}_p: H| = |\lambda_{11}\lambda_{22} \cdots \lambda_{RR}| = p^{-n}. \quad (6)$$

Often we will identify a good basis $h_1, \dots, h_R$ with the R^2 -tuple of co-ordinates (λ_{ij}) .

Proposition 7.1 ([4], Lemma 2.3) *The set $\mathcal{D}$ of good bases $(\lambda_{ij}) \in \mathbb{Z}_p^{R^2}$ is definable.*

For each nonnegative n consider the following:

- (a) the number of index p^n subgroups of Δ ;
- (b) the number of normal index p^n subgroups of Δ ;

- (c) the number of index p^n subgroups A of Δ such that $\widehat{A}_p \simeq \widehat{\Delta}_p$;
- (d) the number of conjugacy classes of index p^n subgroups of Δ ;
- (e) the number of equivalence classes of index p^n subgroups of Δ , where we define $A \sim B$ if $\widehat{A}_p \simeq \widehat{B}_p$.

Parts (a)–(d) of the following result are due to Grunewald, Segal, and Smith [4], Theorem 1.

Proposition 7.2 *Let b_n be as described in any of (a)–(e) above. Then the local zeta function $\sum_{n=0}^{\infty} b_n t^n$ is $\mathbb{Q}$ -rational.*

Proof Consider case (a). Let $\mathcal{D}$ be as in Proposition 7.1. Define $f: \mathcal{D} \rightarrow \mathbb{Q}_p$ by $f(\lambda_{ij}) = |\lambda_{11} \cdots \lambda_{RR}|$. Define an equivalence relation $\mathcal{E}$ on $\mathcal{D}$ as follows: two R -tuples (λ_{ij}) , (μ_{ij}) , representing good bases $h_1, \dots, h_R$ and $k_1, \dots, k_R$ for subgroups H , K respectively, are equivalent if and only if $H = K$. Using Eqn. (6), we see that for each $n \in \mathbb{Z}$, the subset $\{(\lambda_{ij}) \in \mathcal{D} \mid |f(\lambda_{ij})| = p^{-n}\}$ is the union of precisely b_n equivalence classes.

Now $\mathcal{E}$ is definable: it is the subset of $\mathcal{D} \times \mathcal{D}$ given by the conjunction for $1 \leq i, j \leq R$ of the formulae

$$(\exists \sigma_1^{(i)}, \dots, \sigma_R^{(i)} \in \mathbb{Z}_p) \ k_i = h_1^{\sigma_1^{(i)}} \cdots h_R^{\sigma_R^{(i)}}$$

and

$$(\exists \tau_1^{(j)}, \dots, \tau_R^{(j)} \in \mathbb{Z}_p) \ k_j = h_1^{\tau_1^{(j)}} \cdots h_R^{\tau_R^{(j)}},$$

and these become polynomial equations in the λ_{ij} , the μ_{ij} , the σ_i and the τ_j when we write the h_i and k_j in terms of their Mal'cev co-ordinates (Eqn. (5)). We deduce from Theorem 6.2 that $\sum_{n=0}^{\infty} b_n t^n$ is $\mathbb{Q}$ -rational.

The proofs in cases (b) – (e) are similar, modifying the definitions of $\mathcal{D}$ and $\mathcal{E}$ appropriately. For example, in (b) we replace $\mathcal{D}$ by the set $\mathcal{D}^{\trianglelefteq}$ of tuples (λ_{ij}) that define a normal finite-index subgroup H ; a tuple (λ_{ij}) corresponding to a finite index subgroup H belongs to $\mathcal{D}^{\trianglelefteq}$ if and only if it satisfies the formula

$$(\forall g \in \widehat{\Delta}_p)(\forall h \in H)(\exists \nu_1, \dots, \nu_R \in \mathbb{Z}_p) \ ghg^{-1} = h_1^{\nu_1} \cdots h_R^{\nu_R},$$

which is polynomial in the ν_i , the λ_{ij} and the Mal'cev co-ordinates of g and h . In case (d), the equivalence relation is the subset of $\mathcal{D} \times \mathcal{D}$ given by the formula:

there exists $g \in \widehat{\Delta}_p$, there exists $\sigma_i^{(j)}, \tau_i^{(j)} \in \mathbb{Z}_p$ for $1 \leq i, j \leq R$ such that

$$gh_jg^{-1} = k_1^{\sigma_1^{(j)}} \cdots k_R^{\sigma_R^{(j)}} \text{ and } gk_jg^{-1} = h_1^{\tau_1^{(j)}} \cdots h_R^{\tau_R^{(j)}} \text{ for } 1 \leq j \leq R.$$

This is polynomial in the Mal'cev co-ordinates of g and of the h_i and the k_i .

The proof for case (d) is not given explicitly in [4], but the appropriate definable integral can be constructed using the methods in the proof of [dS, Conj. classes], Theorem 1.2; what makes this work is that the equivalence classes are the orbits of a group action. (Note, however, that our method is not directly applicable to [dS, ibid] because the language used there contains symbols for analytic functions.) Case (e) is new; here the equivalence relation does not arise from any obvious group action, and Theorem 6.2 gives a genuinely new way of proving rationality.

Here is another application, to the problem of counting finite p -groups.

Proposition 7.3 (du Sautoy) *Fix positive integers c, d . Let c_n be the number of finite p -groups of order p^n and nilpotency class at most c , generated by at most d elements. Then the power series $\sum_{n=0}^{\infty} c_n t^n$ is a rational function of t .*

Proof Let Δ be the free nilpotent group of class c on d generators. Any finite p -group of order p^n and nilpotency class at most c and generated by at most d elements is a quotient of $\hat{\Delta}_p$ by some normal subgroup of index p^n . Let $\mathcal{D}^{\trianglelefteq}$ and f be as in the proof of Proposition 7.2. Define an equivalence relation $\mathcal{E}$ on $\mathcal{D}^{\trianglelefteq}$ as follows: two R -tuples $(\lambda_{ij}), (\mu_{ij})$, representing good bases $h_1, \dots, h_R$ and $k_1, \dots, k_R$ for subgroups H, K respectively, are equivalent if and only if $\hat{\Delta}_p/H \simeq \hat{\Delta}_p/K$.

The result will follow as in Proposition 7.2 if we can show that $\mathcal{E}$ is definable. We claim that $\mathcal{E} \subset \mathcal{D}^{\trianglelefteq} \times \mathcal{D}^{\trianglelefteq}$ is given by conjunction of the formulae

$$|f(\lambda_{ij})| = |f(\mu_{ij})|$$

and

$$|(\exists b_1, \dots, b_r \in \hat{\Delta}_p)(\forall \nu_1, \dots, \nu_r \in \mathbb{Z}_p) a_1^{\nu_1} \cdots a_R^{\nu_R} \in H \iff b_1^{\nu_1} \cdots b_R^{\nu_R} \in K|. \quad (7)$$

[There is a problem here. E.g., the a_i aren't defined/quantified over.]

To prove this, suppose that $|\hat{\Delta}_p : H| = |\hat{\Delta}_p : K|$. If Eqn. (7) holds then the map $a_i H \mapsto b_i K$ defines an isomorphism from $\hat{\Delta}_p/H$ onto $\hat{\Delta}_p/K$. Conversely,

if g is an isomorphism from $\widehat{\Delta}_p/H$ onto $\widehat{\Delta}_p/K$ then we can choose $b_i \in \widehat{\Delta}_p$ such that $g(a_i H) = b_i K$ for $1 \leq i \leq R$. Then for all $\nu_1, \dots, \nu_r \in \mathbb{Z}$ we have

$$(*) \quad a_1^{\nu_1} \cdots a_R^{\nu_R} \in H \iff b_1^{\nu_1} \cdots b_R^{\nu_R} \in K;$$

since H, K are closed and the group operations are continuous, $(*)$ holds with $\mathbb{Z}$ replaced by $\mathbb{Z}_p$. This proves the claim. The formulae above involve only the definable function f , the p -adic valuation, and polynomials in the Mal'cev co-ordinates and the ν_i , so $\mathcal{E}$ is definable, as required.

Du Sautoy's proof [ref:zeta fns and counting finite gps, Thm 2.2] uses the fact that an isomorphism $\widehat{\Delta}_p/H \rightarrow \widehat{\Delta}_p/K$ lifts to an automorphism of $\widehat{\Delta}_p$, which implies that the equivalence relation $\mathcal{E}$ arises from the action of the group $\text{Aut } \widehat{\Delta}_p$, a compact p -adic analytic group. This allows one to express the power series $\sum_{n=0}^{\infty} c_n t^n$ as a definable p -adic cone integral. Our proof is simpler; on the other hand, it does not yield the information about uniformity in p that follows from the cone integral formulation ([dS, ibid. Thm 3.2]).

8 Twist Isoclasses of Characters of Nilpotent Groups

By a representation of a group G we shall mean a finite-dimensional complex representation, and by a character of G we shall mean the character of such a representation. A character is said to be linear if its degree is one. We write $\langle, \rangle_G$ for the usual inner product of characters of G . If χ is linear then we have

$$\langle \chi \sigma_1, \chi \sigma_2 \rangle_G = \langle \sigma_1, \sigma_2 \rangle_G \quad (8)$$

for all characters σ_1 and σ_2 . If $G' \leq G$ has finite index then we write $\text{Ind}_{G'}^G \cdot$ and $\text{Res}_{G'}^G \cdot$ for the induced character and restriction of a character respectively. For background on representation theory, see [1]. Below when we apply results from the representation theory of finite groups to representations of an infinite group, the representations concerned always factor through finite quotients.

We denote the set of irreducible n -dimensional characters of G by $R_n(G)$. If $N \trianglelefteq G$ then we say the character χ of a representation ρ factors through G/N if ρ factors through G/N (this depends only on χ , not on ρ).

Notation 8.1 We say a character σ of G is *admissible* if σ factors through a finite quotient of G . If p is prime then we say σ is *p-admissible* if σ factors through a finite p -group quotient of G . We write $R_n^{\text{ad}}(G)$ ($R_n^{(p)}(G)$) for the set of admissible (p -admissible) characters in $R_n(G)$. Note that $R_n^{(p)}(G)$ is empty if n is not a p -power ([1], (9.3.2) Proposition).

Given $\sigma_1, \sigma_2 \in R_n(G)$, we follow [8] and say that σ_1 and σ_2 are *twist-equivalent* if $\sigma_1 = \chi\sigma_2$ for some linear character χ of G . Clearly this defines an equivalence relation on $R_n(G)$; we call the equivalence classes *twist isoclasses*.

[Notation: Do we want to say “irreducible twist isoclasses” or just “twist isoclasses”?]

Observation 8.2 Let σ_1, σ_2 be two irreducible degree n characters of G that are twist-equivalent: say $\sigma_2 = \chi\sigma_1$. If $N \trianglelefteq G$ such that σ_1, σ_2 both factor through G/N , then χ also factors through G/N .

If $N_1, N_2 \trianglelefteq G$ have finite (p -power) index then $N_1 \cap N_2$ also has finite (p -power) index. This implies that that when we are working with twist isoclasses in $R_n^{\text{ad}}(G)$ ($R_n^{(p)}(G)$), we need only consider twisting by admissible (p -admissible) linear characters.

Fix a finitely generated nilpotent group Γ . The set $R_n(\Gamma)$ can be given the structure of a quasi-affine complex algebraic variety. Lubotzky and Magid analysed the geometry of this variety and proved the following result [8], [ref?].

Theorem 8.3 *There exists a finite quotient $\Gamma(n)$ of Γ such that every irreducible n -dimensional representation of Γ factors through $\Gamma(n)$ up to twisting. In particular, there are only finitely many twist isoclasses of irreducible n -dimensional characters.*

Thus the number of degree n irreducible twist isoclasses is a finite number a_n .

Theorem 8.4 *The zeta function $\zeta_{\Gamma,p}(t) = \sum_{n=0}^{\infty} a_p^n t^n$ is rational.*

We prove Theorem 8.4 by showing how to parametrise irreducible twist isoclasses in a definable way. The equivalence relation in the parametrisation is not simply the relation of twist-equivalence, which arises from the action of a group — the group of linear characters of Γ — but a more complicated equivalence relation.

The correspondence between index p^n subgroups of Γ and index p^n subgroups of $\widehat{\Gamma}_p$ gives a canonical bijection between $R_{p^n}^{(p)}(\Gamma)$ and $R_{p^n}^{(p)}(\widehat{\Gamma}_p)$, and it is clear that this respects twisting by p -admissible characters.

Lemma 8.5 *For every nonnegative integer n , the sets $R_{p^n}(\Gamma)/(\text{twisting})$ and $R_{p^n}^{(p)}(\widehat{\Gamma}_p)/(\text{twisting})$ are in bijective correspondence with each other.*

Proof It suffices to show that given any $\sigma \in R_{p^n}(\Gamma)$, some twist of σ factors through a finite p -group quotient of Γ . By Theorem 8.3, we can assume that σ factors through some finite quotient F of Γ . Then F , being a finite nilpotent group, is the direct product of its Sylow l -subgroups F_l , where l ranges over all the primes dividing $|F|$ ([ref: Hall], Theorem 2.7). Moreover ([1], (10.33) Theorem), σ is a product of irreducible characters σ_l , where each σ_l is a character of F_l . Since the degree of an irreducible character of a finite group divides the order of the group ([1], (9.3.2) Proposition), all of the σ_l for $l \neq p$ are linear. We may therefore twist σ by a linear character of F to obtain a character that kills F_l for $l \neq p$. The new character factors through F_p , and we are done.

The key idea is that finite p -groups are monomial: that is, every irreducible character is induced from a linear character of some subgroup. We parametrise p -admissible irreducible characters of $\widehat{\Gamma}_p$ by certain pairs (H, χ) , where H is a finite index subgroup of $\widehat{\Gamma}_p$ and χ is a p -admissible linear character of H : to a pair we associate the induced character $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$. We can parametrise these pairs using the theory of good bases for subgroups of $\widehat{\Gamma}_p$, and this description is well-behaved with respect to twisting. Two distinct pairs (H, χ) and (H', χ') may give the same induced character; this gives rise to a definable equivalence relation on the set of pairs.

If ψ is a character of H then we denote by $g.\psi$ the character of $g.H = gHg^{-1}$ defined by $(g.\psi)(ghg^{-1}) = \psi(g)$.

Lemma 8.6 (a) *Let $\sigma \in R_{p^n}^{(p)}(\widehat{\Gamma}_p)$. Then there exists $H \leq \widehat{\Gamma}_p$, such that $|\widehat{\Gamma}_p : H| = p^n$, together with a p -admissible linear character χ of H such that $\sigma = \text{Ind}_H^{\widehat{\Gamma}_p} \chi$.*

(b) *Let H be a p -power index subgroup of $\widehat{\Gamma}_p$ and let χ be a p -admissible linear character of H . Then $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ is a p -admissible character of $\widehat{\Gamma}_p$, and $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ is irreducible if and only if for all $g \in \widehat{\Gamma}_p - H$, $\text{Res}_{g.H \cap H}^{g.H} g.\chi \neq \text{Res}_{g.H \cap H}^H \chi$.*

Moreover, if ψ is a p -admissible linear character of $\widehat{\Gamma}_p$ and $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ is irreducible then $\text{Ind}_H^{\widehat{\Gamma}_p} \left(\left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \chi \right) = \psi \text{Ind}_H^{\widehat{\Gamma}_p} \chi$.

(c) Let $H, H' \leq \widehat{\Gamma}_p$ have index p^n , and let χ, χ' be p -admissible linear characters of H, H' respectively such that $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ and $\text{Ind}_{H'}^{\widehat{\Gamma}_p} \chi'$ are irreducible. Then $\text{Ind}_H^{\widehat{\Gamma}_p} \chi = \text{Ind}_{H'}^{\widehat{\Gamma}_p} \chi'$ if and only if there exists $g \in \widehat{\Gamma}_p$ such that $\text{Res}_{g.H \cap H'}^{g.H} g \cdot \chi = \text{Res}_{g.H \cap H'}^{H'} \chi'$.

Proof (a) Since σ is p -admissible, it factors through some finite p -group F . Since finite p -groups are monomial ([1], (11.3) Theorem), there exists $L \leq F$ of index p^n and a linear character χ of L such that σ — regarded as a character of F — equals $\text{Ind}_L^F \chi$. Let H be the preimage of L under the canonical projection $\widehat{\Gamma}_p \rightarrow F$. Regarding χ as a character of H , it is easily checked that $|\widehat{\Gamma}_p : H| = p^n$ and $\sigma = \text{Ind}_H^{\widehat{\Gamma}_p} \chi$.

(b) Since χ is p -admissible, the kernel K of χ has p -power index in $\widehat{\Gamma}_p$, so K contains a p -power index subgroup N such that $N \trianglelefteq \widehat{\Gamma}_p$. Clearly $N \leq \ker(\text{Ind}_H^{\widehat{\Gamma}_p} \chi)$, so $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ is p -admissible. The irreducibility criterion follows immediately from [1], (10.25) Theorem. Using Frobenius Reciprocity ([7], Chapter XVIII, Theorem 6.1), we have

$$\begin{aligned}
& \left\langle \text{Ind}_H^{\widehat{\Gamma}_p} \left(\left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \chi \right), \psi \text{Ind}_H^{\widehat{\Gamma}_p} \chi \right\rangle_{\widehat{\Gamma}_p} \\
&= \left\langle \left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \chi, \text{Res}_H^{\widehat{\Gamma}_p} \left(\psi \text{Ind}_H^{\widehat{\Gamma}_p} \chi \right) \right\rangle_H \\
&= \left\langle \left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \chi, \left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \text{Res}_H^{\widehat{\Gamma}_p} \left(\text{Ind}_H^{\widehat{\Gamma}_p} \chi \right) \right\rangle_H \\
&= \left\langle \chi, \text{Res}_H^{\widehat{\Gamma}_p} \left(\text{Ind}_H^{\widehat{\Gamma}_p} \chi \right) \right\rangle_H \text{ by Eqn. (8)} \\
&= \left\langle \text{Ind}_H^{\widehat{\Gamma}_p} \chi, \text{Ind}_H^{\widehat{\Gamma}_p} \chi \right\rangle_{\widehat{\Gamma}_p} \\
&= 1.
\end{aligned}$$

Now $\psi \text{Ind}_H^{\widehat{\Gamma}_p} \chi$ is irreducible, because $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ is, and the degrees of $\text{Ind}_H^{\widehat{\Gamma}_p} \left(\left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \chi \right)$ and $\psi \text{Ind}_H^{\widehat{\Gamma}_p} \chi$ are equal. We deduce that $\psi \text{Ind}_H^{\widehat{\Gamma}_p} \chi = \text{Ind}_H^{\widehat{\Gamma}_p} \left(\left(\text{Res}_H^{\widehat{\Gamma}_p} \psi \right) \chi \right)$.

(c) The Mackey Subgroup Theorem [1], (10.13) Theorem gives

$$\text{Res}_{H'}^{\widehat{\Gamma}_p} \left(\text{Ind}_H^{\widehat{\Gamma}_p} \chi \right) = \sum_{\overline{g} \in H' \backslash \widehat{\Gamma}_p / H} \text{Ind}_{g.H \cap H'}^{H'} \left(\text{Res}_{g.H \cap H'}^{g.H} g \cdot \chi \right). \quad (9)$$

Here the sum is over a set of double coset representatives g for $H' \backslash \widehat{\Gamma}_p / H$ (the characters on the RHS of the formula are independent of choice of representative). Since $\text{Ind}_H^{\widehat{\Gamma}_p} \chi$ and $\text{Ind}_{H'}^{\widehat{\Gamma}_p} \chi'$ are irreducible, they are distinct if and only if their inner product is zero. We have

$$\begin{aligned} & \left\langle \text{Ind}_H^{\widehat{\Gamma}_p} \chi, \text{Ind}_{H'}^{\widehat{\Gamma}_p} \chi' \right\rangle_{\widehat{\Gamma}_p} \\ &= \left\langle \text{Res}_{H'}^{\widehat{\Gamma}_p} (\text{Ind}_H^{\widehat{\Gamma}_p} \chi), \chi' \right\rangle_{H'} \text{ by Frobenius Reciprocity} \\ &= \sum_{\overline{g} \in H' \backslash \widehat{\Gamma}_p / H} \left\langle \text{Ind}_{g.H \cap H'}^{H'} \left(\text{Res}_{g.H \cap H'}^{g.H} g \cdot \chi \right), \chi' \right\rangle_{H'} \text{ by the Mackey Subgroup Theorem} \\ &= \sum_{\overline{g} \in H' \backslash \widehat{\Gamma}_p / H} \left\langle \text{Res}_{g.H \cap H'}^{g.H} g \cdot \chi, \text{Res}_{g.H \cap H'}^{H'} \chi' \right\rangle_{g.H \cap H'} \text{ by Frobenius Reciprocity.} \end{aligned}$$

This vanishes if and only if each of the summands vanishes, which happens if and only if $\text{Res}_{g.H \cap H'}^{g.H} g \cdot \chi \neq \text{Res}_{g.H \cap H'}^{H'} \chi'$ for every g , since the characters concerned are linear. The result follows.

Write Γ as a quotient Δ/Θ of a finitely generated torsion-free nilpotent group Δ : for example, we may take Δ to be the free class c nilpotent group on N generators for appropriate N and c . Let $\pi: \Delta \rightarrow \Gamma$ be the canonical projection, and let $i: \Theta \rightarrow \Delta$ be inclusion. Let $\widehat{\Delta}_p, \widehat{\Theta}_p$ be the pro- p completions of Δ, Θ respectively. Then π (resp. i) extends to a continuous homomorphism $\widehat{\pi}_p: \widehat{\Delta}_p \rightarrow \widehat{\Gamma}_p$ (resp. $\widehat{i}_p: \widehat{\Theta}_p \rightarrow \widehat{\Delta}_p$), and the three groups $\widehat{i}_p(\widehat{\Theta}_p)$, $\ker \widehat{\pi}_p$, and the closure of Θ in $\widehat{\Delta}_p$ all coincide (compare [3], Exercise 21 of Chapter 1). (Because Δ is finitely generated nilpotent, it can in fact be shown that $\widehat{i}_p$ is injective, and hence an isomorphism onto its image.) Clearly p -admissible representations of $\widehat{\Gamma}_p$ correspond bijectively to p -admissible representations of $\widehat{\Delta}_p$ that kill $\ker \widehat{\pi}_p$. By [ref: Hall], Lemma 1.9, Θ is finitely generated, so we can choose a Mal'cev basis $\theta_1, \dots, \theta_s$ for Θ . We identify the θ_i with their images in $\widehat{\Delta}_p$.

Let μ_{p^n} be the group of all complex p^n th roots of unity, and let μ_{p^∞} be the group of all complex p -power roots of unity.

Lemma 8.7 *The groups μ_{p^∞} and $\mathbb{Q}_p/\mathbb{Z}_p$ are isomorphic.*

Proof Let $p^{-\infty}\mathbb{Z} \leq \mathbb{Q}$ be the group of rational numbers of the form np^{-r} for $n \in \mathbb{Z}$ and r a nonnegative integer. Then $p^{-\infty}\mathbb{Z} \cap \mathbb{Z}_p = \mathbb{Z}$ and $\mathbb{Z}_p p^{-\infty}\mathbb{Z} = \mathbb{Q}_p$, so $\mathbb{Q}_p/\mathbb{Z}_p \simeq p^{-\infty}\mathbb{Z}/\mathbb{Z}$, by one of the standard group isomorphism theorems. The map $q \mapsto e^{2\pi i q}$ gives an isomorphism from $p^{-\infty}\mathbb{Z}/\mathbb{Z}$ to μ_{p^∞} .

Let $\Phi: \mu_{p^\infty} \rightarrow \mathbb{Q}_p/\mathbb{Z}_p$ be the isomorphism described above. Any p -admissible linear character of a pro- p group takes its values in μ_{p^∞} , so we use Φ to identify p -admissible linear characters with p -admissible homomorphisms to $\mathbb{Q}_p/\mathbb{Z}_p$.

Lemma 8.8 *Let $\mathcal{D} \subset \mathbb{Z}_p^{R^2} \times \mathbb{Q}_p^R$ be the set of tuples (λ_{ij}, y_k) , where $1 \leq i, j \leq R$ and $1 \leq k \leq R$, satisfying the following conditions:*

- (a) *the λ_{ij} form a good basis $h_1, \dots, h_R$ for some finite index subgroup H of $\hat{\Delta}_p$;*
- (b) *the prescription $h_i \mapsto y_i \bmod \mathbb{Z}_p$ gives a well-defined p -admissible homomorphism $\chi: H \rightarrow \mathbb{Q}_p/\mathbb{Z}_p$ that kills $H \cap \ker \hat{\pi}_p$;*
- (c) *the induced character $\text{Ind}_H^{\hat{\Delta}_p} \chi$ is irreducible.*

Then $\mathcal{D}$ is definable. Moreover, $\text{Ind}_H^{\hat{\Delta}_p} \chi$ is a p -admissible character of $\hat{\Gamma}_p$, and every p -admissible irreducible character of $\hat{\Gamma}_p$ arises in this way. [Want $\pi_p \circ \dots$ for this to make sense.]

Notation 8.9 Given $(\lambda_{ij}, y_k) \in \mathcal{D}$, we write $\Psi(\lambda_{ij}, y_k)$ for the pair (H, χ) . Since the h_i generate H topologically, the p -admissible homomorphism χ defined by the y_i is unique.

Proof Condition (a) is definable, by Proposition 7.1. Given that (a) holds, we claim that (b) holds if and only if there exists an R^2 -tuple (μ_{ij}) such that:

- (i) (μ_{ij}) defines a good basis $k_1, \dots, k_R$ for a finite index subgroup K of $\hat{\Delta}_p$;
- (ii) $K \leq H$;
- (iii) $H \cap \ker \hat{\pi}_p \subset K$;
- (iv) there exist $y \in \mathbb{Q}_p$, $r_1, \dots, r_R \in \mathbb{Z}_p$, $h \in H$ such that $|y| = |H/K|$ and for every i we have $\overline{h^{r_i}} = \overline{h_i}$ and $r_i y = y_i \bmod \mathbb{Z}_p$. (Here $\overline{x}$ denotes the image of $x \in H$ under the canonical projection $H \rightarrow H/K$.)

To see this, note that if (b) holds then $\ker \chi$ is a finite index subgroup of H which satisfies (ii) and (iii). Take (μ_{ij}) to be any tuple defining a good basis

for K . Then H/K , being a finite subgroup of $\mathbb{Q}_p/\mathbb{Z}_p$, is cyclic, so choose $h \in H$ that generates H/K and choose $y \in \mathbb{Q}_p$ such that $\chi(h) = y \bmod \mathbb{Z}_p$. We can choose $r_1, \dots, r_R \in \mathbb{Z}$ such that $h_i = h^{r_i}$ for each i , and it is easily checked that (iv) holds.

Conversely, suppose there exists a tuple (μ_{ij}) satisfying (i)–(iv). The map $\mathbb{Z}_p \rightarrow H$, $\lambda \mapsto h^\lambda$ is continuous because it is polynomial with respect to the Mal'cev co-ordinates, so there exists an open neighbourhood U of 0 in $\mathbb{Z}_p$ such that $h^\lambda \in K$ for all $\lambda \in U$. Since $\mathbb{Z}$ is dense in $\mathbb{Z}_p$, we may therefore find $n_1, \dots, n_R \in \mathbb{Z}$ such that $\overline{h_i} = \overline{h^{n_i}}$ for each i .

We have a monomorphism $\beta: H/K \rightarrow \mathbb{Q}_p/\mathbb{Z}_p$ given by $\beta(\overline{h^n}) = ny \bmod \mathbb{Z}_p$. Let χ be the composition $H \rightarrow H/K \xrightarrow{\beta} \mathbb{Q}_p/\mathbb{Z}_p$. Condition (iii) implies that χ kills $H \cap \ker \hat{\pi}_p$. The canonical projection $H \rightarrow H/K$ is continuous ([3], 1.2 Proposition), so we have $\chi(h^\lambda) = \lambda y \bmod \mathbb{Z}_p$ for every $\lambda \in \mathbb{Z}_p$. Condition (iv) implies that $\chi(h_i) = y_i \bmod \mathbb{Z}_p$ for every i , as required.

Now condition (i) is definable, by Proposition 7.1. Condition (iii) can be expressed as

$$\begin{aligned} (\forall \nu_1, \dots, \nu_s \in \mathbb{Z}_p) [(\exists \sigma_1, \dots, \sigma_s \in \mathbb{Z}_p) \theta_1^{\nu_1} \cdots \theta_s^{\nu_s} = h_1^{\sigma_1} \cdots h_R^{\sigma_R}] \\ \Rightarrow (\exists \tau_1, \dots, \tau_s \in \mathbb{Z}_p) \theta_1^{\nu_1} \cdots \theta_s^{\nu_s} = k_1^{\tau_1} \cdots k_R^{\tau_R}. \end{aligned} \quad (10)$$

Eqn. 10 can be expressed in terms of polynomials in the λ_{ij} , the μ_{ij} , the ν_k , the σ_k and the τ_k (note that the θ_k are fixed elements of $\hat{\Delta}_p$).

Similar arguments show that (ii) and (iv) are also definable conditions. In (iv), note that the conditions $\overline{h^{r_i}} = \overline{h_i}$ imply by the argument above that h is a generator for H/K , so the condition $|y| = |H/K|$ can be expressed as $(h^y \in K) \wedge (h^{p^{-1}y} \notin K)$.

By Lemma 8.6 (b), irreducibility of the induced character can be written as:

$$(\forall g \in \hat{\Delta}_p - H)(\exists h \in H) ghg^{-1} \in H \text{ and } \chi(ghg^{-1}) \neq \chi(h).$$

Writing this in terms of the Mal'cev co-ordinates, we see that (c) is a definable condition.

By Lemma 8.6 (a), any p -admissible irreducible character σ of $\hat{\Gamma}_p$ is of the form $\text{Ind}_L^{\hat{\Gamma}_p} \chi$ for some finite index subgroup L of $\hat{\Gamma}_p$ and some p -admissible linear character χ of L . Let H be the preimage of L under the canonical projection $\hat{\Delta}_p \rightarrow \hat{\Gamma}_p$. Regarding σ, χ as representations of $\hat{\Delta}_p, H$ respectively, it is easily checked that $\sigma = \text{Ind}_H^{\hat{\Delta}_p} \chi$. Choose (λ_{ij}) defining a good basis

$h_1, \dots, h_R$ for H , and choose y_k such that $\chi(h_i) = y_i \bmod \mathbb{Z}_p$. The above argument shows that $(\lambda_{ij}, y_k) \in \mathcal{D}$. This completes the proof.

Define $f: \mathcal{D} \rightarrow \mathbb{Z}_p$ by $f(\lambda_{ij}, y_k) = |\lambda_{11} \cdots \lambda_{RR}|$. Define an equivalence relation $\mathcal{E}$ on $\mathcal{D}$ by $(\lambda_{ij}, y_k) \sim (\lambda'_{ij}, y'_k)$ if $\text{Ind}_{\widehat{H}}^{\widehat{\Delta}_p} \chi$ and $\text{Ind}_{\widehat{H}'}^{\widehat{\Delta}_p} \chi'$ are twist-equivalent, where $(H, \chi) = \Psi(\lambda_{ij}, y_k)$ and $(H', \chi') = \Psi(\lambda'_{ij}, y'_k)$. The degree of $\text{Ind}_{\widehat{H}}^{\widehat{\Delta}_p} \chi$ equals $f(\lambda_{ij}, y_k)^{-1}$, and likewise for (λ'_{ij}, y'_k) , so if $(\lambda_{ij}, y_k) \sim (\lambda'_{ij}, y'_k)$ then $f(\lambda_{ij}, y_k) = f(\lambda'_{ij}, y'_k)$.

For each nonnegative integer n , let $\mathcal{D}_n = \{(\lambda_{ij}, y_k) \in \mathcal{D} \mid f(\lambda_{ij}, y_k) = p^{-n}\}$. It follows from Lemma 8.8 and the definition of $\mathcal{E}$ that $\mathcal{D}_n$ is the union of precisely a_{p^n} $\mathcal{E}$ -equivalence classes (note that if one representation of $\widehat{\Gamma}_p$ is the twist of another by some linear character ψ of $\widehat{\Delta}_p$ then ψ is automatically a character of $\widehat{\Gamma}_p$, by Observation 8.2). We may regard $\mathcal{E}$ as a family $(\mathcal{E}_n)_{n \in \mathbb{Z}}$ of equivalence relations on the definable family of sets $(\mathcal{D}_n)_{n \in \mathbb{Z}}$. To complete the proof of Theorem 8.4, it suffices by Theorem 6.2 to prove the following result.

Proposition 8.10 *The equivalence relation $\mathcal{E}$ is definable.*

Proof Let $\mathcal{D}' \subset \mathbb{Q}_p^R$ be the set of R -tuples $(z_1, \dots, z_R)$ such that the prescription $a_i \mapsto z_i \bmod \mathbb{Z}_p$ gives a well-defined p -admissible linear character of $\widehat{\Delta}_p$ that kills $\ker \widehat{\pi}_p$. We denote this character by $\Xi(z_1, \dots, z_R)$ (or just $\Xi(z_k)$). Similar arguments to those in the proof of Lemma 8.8 show that $\mathcal{D}'$ is definable. If $(z_1, \dots, z_R) \in \mathcal{D}'$ and $(H, \chi) = \Psi(\lambda_{ij}, y_k)$ then $\text{Res}_{\widehat{H}}^{\widehat{\Delta}_p} \Xi(z_k) \chi = \Psi(\lambda_{ij}, y_k + z_k)$. Applying Lemma 8.6 (b) and (c), we see that $(\lambda_{ij}, y_k) \sim (\lambda'_{ij}, y'_k)$ if and only if

$$\begin{aligned} (\exists (z_1, \dots, z_R) \in \mathcal{D}') (\exists g \in \widehat{\Delta}_p) (\forall h \in H) ghg^{-1} \in H' \\ \Rightarrow \left(\text{Res}_{\widehat{H}}^{\widehat{\Delta}_p} \Xi(z_k) \chi \right) (h) = \chi'(ghg^{-1}). \end{aligned}$$

Writing this in terms of the Mal'cev co-ordinates, we obtain an equation involving $\mathcal{D}'$ and p -adic norms of polynomials in the λ_{ij} , the y_k , the λ'_{ij} , the y'_k , the z_k , and the Mal'cev co-ordinates of g and h . We deduce that $\mathcal{E}$ is definable, as required.

Remark 8.11 It is not hard to see that the sets $\mathcal{D}, \mathcal{E}$ are actually 0-definable, for all the polynomials that arise have coefficients in $\mathbb{Z}$. For example, since each θ_i belongs to Θ , its Mal'cev co-ordinates with respect to the a_i are integers.

Example 8.12 Let Γ be the Heisenberg group $\langle a, b, c \mid [a, b] = c, [a, c] = [b, c] = 1 \rangle$. Nunley and Magid [10] explicitly calculated the irreducible twist isoclasses of Γ and showed that

$$\zeta_{\Gamma,p}(t) = 1 + \sum_{n=1}^{\infty} (p-1)p^{n-1}t^n.$$

The formula for the sum of a geometric progression gives

$$\zeta_{\Gamma,p}(t) = \frac{1-t}{1-pt}.$$

References

- [1] Charles W. Curtis and Irving Reiner, *Methods of representation theory. Vol. I*, John Wiley & Sons Inc, New York, 1981, With applications to finite groups and orders, Pure and Applied Mathematics, A Wiley-Interscience Publication.
- [2] Jan Denef, *Arithmetic and geometric applications of quantifier elimination for valued fields*, Model theory, algebra, and geometry, Math. Sci. Res. Inst. Publ., vol. 39, Cambridge Univ. Press, Cambridge, 2000, pp. 173–198.
- [3] J. D. Dixon, M. P. F. du Sautoy, A. Mann, and D. Segal, *Analytic pro- p groups*, second ed., Cambridge Studies in Advanced Mathematics, vol. 61, Cambridge University Press, Cambridge, 1999.
- [4] F. J. Grunewald, D. Segal, and G. C. Smith, *Subgroups of finite index in nilpotent groups*, Invent. Math. **93** (1988), no. 1, 185–223.
- [5] Deirdre Haskell, Ehud Hrushovski, and Dugald Macpherson, *Definable sets in algebraically closed valued fields. part I: elimination of imaginaries*, Preprint, 2004.
- [6] ———, *Definable sets in algebraically closed valued fields. part II: stable domination and independence*, Preprint, 2004.
- [7] Serge Lang, *Algebra*, third ed., Addison-Wesley, Reading, Massachusetts, 1993.

- [8] Alexander Lubotzky and Andy R. Magid, *Varieties of representations of finitely generated groups*, Mem. Amer. Math. Soc. **58** (1985), no. 336, xi+117.
- [9] Alexander Lubotzky and Benjamin Martin, *Polynomial representation growth and the congruence subgroup problem*, Israel J. Math., to appear, 2003.
- [10] Charles Nunley and Andy Magid, *Simple representations of the integral Heisenberg group*, Classical groups and related topics (Beijing, 1987), Contemp. Math., vol. 82, Amer. Math. Soc., Providence, RI, 1989, pp. 89–96.
- [11] Lou van den Dries, *Algebraic theories with definable Skolem functions*, J. Symbolic Logic **49** (1984), no. 2, 625–629.